



**Survey and Nutritive Scanning of Low-cost
Feedstuffs and Energy Requirement of
Indian Major Carp, *Labeo rohita* (Ham)**

DISSERTATION

**SUBMITTED IN PARTIAL FULFILMENT OF THE REQUIREMENTS
FOR THE AWARD OF THE DEGREE OF**

Master of Philosophy

IN

ZOOLOGY

BY

Md. Abul Hassan

**FISH NUTRITION RESEARCH LABORATORY
DEPARTMENT OF ZOOLOGY
ALIGARH MUSLIM UNIVERSITY
ALIGARH (INDIA)**

1989



DS1487



DEPARTMENT OF ZOOLOGY
ALIGARH MUSLIM UNIVERSITY
ALIGARH—202 002, U.P.
(INDIA)

Phones : [University : 285
[Public : 6646

Sections :


1. ENTOMOLOGY
2. PARASITOLOGY
3. ICHTHYOLOGY & FISHERIES
4. AGRICULTURAL NEMATOLOGY
5. GENETICS

Ref. No.

Date 8-11-1989

This is to certify that the dissertation entitled "Survey and Nutritive Scanning of Low-cost Feedstuffs and Energy Requirement of Indian Major Carp, Labeo rohita (Ham.) has been completed under my supervision by Md. Abul Hassan. This work is original and has been independently persued by the candidate.

I permit the candidate to submit the dissertation in partial fulfilment for the award of the degree of Master of Philosophy in Zoology of the Aligarh Muslim University, Aligarh.


A.K. Jafri
M.Sc., Ph.D., F.N.A.Sc.
Professor.

C O N T E N T S

	Page
ACKNOWLEDGEMENTS	
GENERAL INTRODUCTION	1
CHAPTERS	
I. Nutrient composition and energy density of low - cost feedstuffs for their poss- ible use in practical fish feed.	7
II. Energy requirement of the Indian major carp, <u>Labeo rohita</u> (Ham).	24
REFERENCES	38

ACKNOWLEDGEMENTS

I am indebted beyond measure to Prof. A.K.Jafri, for his inspiration, constructive criticism and overall guidance during the course of the present study.

I offer my sincere thanks to the Chairman, Department of Zoology, Aligarh Muslim University, Aligarh, for providing the necessary laboratory facilities.

I am also indebted to Prof. John E. Halver, School of Fisheries, University of Washington, Seattle, USA, for his valuable suggestions and criticism.

All my colleagues, Messers Mukhtar A. Khan, M.F. Anwar, Erfanullah and Md. Nasiruddin, deserve thanks for their cooperation and help.

The financial assistance rendered by US Department of Agriculture is gratefully acknowledged.

GENERAL INTRODUCTION

The increasing demand for high quality low-cost animal protein has generated greater interest to develop and expand aquaculture all over the world. It is realised that scientific exploitation of cultivable water bodies can increase fish production manifold, and it has been predicted that, by the year 2000 AD, aquaculture could produce at least 50 million tons of animal protein, if proper research and development measures are undertaken (TAC, 1973) against a production of 10.5 million tons of fish and shellfish recorded through culture in 1983 (Pillay, 1986).

In India, until recently, extensive or traditional type of fish culture, with production as low as 1 ton/ha/yr or even less, was practiced by most fish farmers. Although low-density culture with minimum inputs and low production per unit area often proved more economical under certain situations than the intensive farming, involving the rearing of dense population and heavy inputs (Pillay, 1976), scientific culture practice and efficient management could lead to a substantial increase in fish production. Production rate as high as 10 tons/ha/yr has been achieved in static earthen freshwater carp culture ponds in India through optimum stocking, fertilization and supplementary feeding (Sinha, 1988).

Fish do require all the vital elements for their growth, reproduction, health and other normal physiological functions. A deficiency of one or more of the essential nutrients results in a reduced rate of performance, disease or even death. In different forms of aquaculture system, fish gather these nutrients from natural aquatic organisms or from the formulated feeds provided in the form of supplementary or whole feed. However, in the farming of these animals under controlled environment, unlike their natural habitat, feeding of the stocked population with nutritionally balanced and quality diets is of critical importance not only to promote their optimal biological and physiological functions but also to raise the production.

In intensive and semi-intensive fish culture systems, based on high stock density, feed is thus the major input, constituting 50-60% of the total operational expenditure. To minimise this input, it is almost essential to develop cost-effective and nutritionally adequate feed, using less expensive locally available ingredients (Pillay, 1986). Such ingredients generally include byproducts of grain/oil industry, fish processing industries, meat processing industries, slaughter house wastes, droppings or wastes of poultry, duckery, dairy and

piggery, etc. Several workers have attempted to scan such ingredients for their nutritive richness in order to incorporate these in low-cost fish diets (Morrison, 1961; Lakshmanan et al., 1967; Chakrabarty et al., 1973; Jhingran 1983, Ensminger and Olentine, Jr, 1980 and NAS-NRC, 1981, 1983).

Fish nutritionists in the past have generally given priority to quantifying the requirements for protein, minerals and vitamins (Halver, 1979; Cowey and Sargent, 1972, 1979; Castell, 1979; Lall, 1979; Nose and Arai, 1979; Ogino, 1979; Stickney, 1979; NAS-NRC, 1981, 1983; Jauncey, 1982; Watanabe, 1982; Ketola, 1982; Millikin, 1982; Takeuchi, et al., 1983; Cowey and Luquet, 1983; Lovell, 1984a; Robinson, 1984; Cho et al., 1985; Kanazawa, 1985; Halver, 1985; Watanabe, 1985; Wilson and Halver, 1986). Although levels and sources of energy in fish ration can significantly affect fish growth, relatively little information is available on the energy requirement of fish.

A factor considered equally important in the diet formulation is the estimation of fibre content. Although non-energy component of diet, the importance of fibre cannot be overlooked in nutritional studies. Chemically, it represents mixtures of cellulose,

hemicellulose, lignin, pentosan and other generally undigestible fractions in the feed. Amongst many functions, it provides increased surface area of substrate to be acted upon by digestive juice, helps in uniform distribution of different nutrients, and facilitates in easy and smooth movement of food material through the alimentary canal, thereby helping in easy defacation and proper digestion. The level of fibre present in the diet has thus a direct role in digestion, absorption and growth efficiency of the animal. Several workers in the past have emphasized the importance of fibre inclusion in artificial fish feed (Dupree and Sneed, 1966; Leary and Lovell, 1975, Stickney and Shumway, 1974; Prejs and Blaszezyk, 1977; and Sumagaysay and Chiu, 1989). Feedstuffs of plant origin contain varied amounts of fibre. Inclusion of such feedstuffs in practical fish diet necessitates evaluation of their fibre content so as to avoid excessive levels of fibre in the diet, mainly because higher levels of fibre may inhibit feed intake, increase fecal waste production, and consequently pollute the water.

Data available on the proximate composition of feedstuffs reported earlier from this laboratory (Niamat, 1982) was neither based on guidelines of the International Network of Feed Information Centre

(INFIC) nor took into account level of fibre content in plant origin feedstuffs and chitin in silkworm pupae. Similarly, the energy values of the feedstuffs reported in the earlier study was based on conventional energy equivalents.

The work presented here was undertaken as part of a research program on the nutrition and diet development of cultivable finfish species. Since formulating rations for feedout or growth increment purposes generally entails consideration of availability, and the biochemical and nutritional features of the ingredients, the study included an assessment of the gross nutrient composition and energy density of locally available feedstuffs.

Besides, fibre has been quantified in the plant origin feedstuffs and chitin in the silkworm pupae. Unlike the earlier study, energy density in the feedstuffs was evaluated through direct combustion in a calorimeter. Additional information relating to the plant origin feedstuffs, as required by INFIC for retrieval purposes, have also been collected. These included details such as the season of harvest, method of preparation in the field, storage condition, types of soil in which grown, fertilizers used and type of farming, etc. The energy requirement for optimum

growth was assessed in the Indian major carp, Labeo rohita (Ham). This is a fast growing, easily cultivable finfish species, and owing to its compatibility with other fish species, highly preferred for cultivation. Although the culture technology of Indian major carps has been perfected in many respects, assessment of nutritional needs of these fishes, leading to the development of complete artificial feed for their rearing, is an aspect which still remains to be precisely worked out to revolutionize the production.

The findings have been outlined under two separate chapters. Chapter I includes findings on the proximate composition and energy density of feedstuffs, Chapter II deals with the energy requirement of L. rohita.

CHAPTER - I

NUTRIENT COMPOSITION AND ENERGY DENSITY OF LOW-COST FEEDSTUFFS FOR THEIR POSSIBLE USE IN PRACTICAL FISH FEED

INTRODUCTION

With the intensification of fish culture operation in many countries of the world, including India, the search for suitable feedstuffs to develop nutritive and economical feed mixtures, based on less expensive locally available feed ingredients, is being increasingly felt (FAO/UNDP, 1979; Halver, 1982), since these contribute greatly to the commercial success of the culture operation. Research efforts are, therefore, directed towards scanning newer and/or economical feed ingredients for aquaculturally important fish species. Aquatic weeds, grasses and leaves have been tested for their suitability as fish feed ingredients (Hajra, 1987; Naskar and Sen, 1987; Pappathi and Paul Raj, 1987; Patra et al., 1989; Cagauan and Pullin, 1989; Naegel, 1989; Paul Raj, 1989; Halinge et al., 1989; Chiayvareesajja et al., 1989 and Klinnavee et al., 1989). Brans, polishing and oil cakes have been tested for their efficacy in practical fish diets (Wee and Little, 1989; Chattopadhyay and Konar, 1987;

Mohanty et al., 1987; and Maria et al., 1987). Agrobased waste products and livestock wastes also find use in the artificial feed for rearing different fish species (Nandesha et al., 1988, and Toor et al., 1988). Use of trash fish, blood meal, maggots and pupae of insects has also been attempted in artificial feed (Otubusin 1987; Nandesha et al., 1989; and Abalos et al., 1989). Efficacy, utilization and level of inclusion of different kinds of organic manure, biogas slurry and slaughter house offal in fish feed formulation were looked into by several workers (Kalyani and Shetty, 1987; Rao , 1987 ; and Haque, 1988). Growth studies on tilapia using single cell protein (Spirulina) has been made by Chow and Woo (1989). Recently, Bhuiyan et al (1989), keeping in view the intensity of aquaculture practice in Bangladesh, conducted an exhaustive survey to identify potential fish feed, based on their availability, price and primary nutritional value.

The present study reports the nutritive richness of locally available ingredients, screened for their proximate composition and energy density, which could find use in developing artificial fish feed.

MATERIALS AND METHODS

The feedstuffs collected from different sources were processed for assessment of their proximate composition and energy contents using standard techniques (AOAC, 1984). Five replicates were taken for each sample.

I. Estimation of water content

For the determination of moisture content each sample was thoroughly grinded to powder. Usually, 2-5 g of the powdered sample was taken in a pre-weighed vitreosil crucible and placed in an oven at 100°C for 20-22 hours. The crucible containing the dried sample was cooled to room temperature in a desiccator and reweighed to ensure that the sample had become completely dried, the entire process was repeated till a constant weight was obtained. The loss in weight gave an index of water from which the percentage of water was calculated. For fresh leafy ingredients the sample were first sun-dried, grinded to fine powder and preserved in polythene bags. This powder was then used for moisture estimation on dry weight basis.

II. Estimation of ash content

A known quantity (1-5 g) of powdered sample was taken in a pre-weighed vitreosil crucible, dried in an electric oven at 100°C and ignited in muffle furnace at 650°C till the sample became carbon-free and completely white. The crucible was cooled in a desiccator and reweighed to estimate the quantity of ash. The result was expressed as percentage on dry weight basis.

III. Estimation of crude fat

Crude fat in different feedstuffs was extracted with petroleum ether (B.P. 40-60°C) using the continuous soxhlet extraction technique. A weighed quantity (5 g) of finely powdered and oven dried (100°C) sample was taken in Whatman fat extraction thimble, plugged with cotton and introduced into the soxhlet. A clean dry receiver flask was weighed and fitted to the soxhlet. The extraction was carried out for about 10-12 hours. At the end of extraction, the solvent was recollected and the flask placed in an oven for few hours for complete removal of solvent traces. The flask was cooled in a desiccator and reweighed. The increase in weight of the flask gave

the quantity of crude fat extracted from the known weight of the sample. The result was expressed as percent on dry weight basis.

IV. Estimation of crude protein

The estimation of crude protein in various samples was made by a slight modification of Wong's micro-kjeldahl method as adopted by Jafri et al. (1964). 0.1 g sample was digested in 5 ml of 1:1 sulphuric acid and heated till fumes disappeared. 0.5 ml saturated potassium persulphate was then added to oxidize the digesting mixture. The digestion was continued till the solution in the Kjeldahl flask became water clear, indicating that all the nitrogenous materials present in the sample has been converted into ammonium sulphate. The solution at this stage was transferred to volumetric flask and diluted to 50 ml. An aliquot (0.5 ml) of this solution was then directly nesslerized by adding Bock and Bendict Nessler's reagent (Oser, 1965). The solution was kept at room temperature for 10 minute for complete colour development and the intensity of colour read on Milton Ray splitbeam spectronic 1001 spectrophotometer at 480 mμ wave length. The intensity of colour developed was proportional to the amount of ammonium sulphate cont-

ained in the solution. The optical density obtained for the sample was read off against a standard calibration curve prepared by taking readings of a series of different dilutions containing known amount of nitrogen. This method gave a direct reading of the amount of total nitrogen present in the sample. The quantity of total nitrogen was multiplied with the conventional protein factor (6.25) to calculate the crude protein content. The values were calculated as percentage on dry weight basis.

V. Estimation of gross energy content

The estimation of gross energy in the feedstuffs was carried out through direct calorimetry. As a routine, five samples were tested for each ingredient. In order to ensure that the test sample is truly representative of the bulk of the sample material, standard sampling, grinding, mixing and subdividing procedures were followed. The sample was dried in hot air oven at 100°C for 24 hour. A weighed portion of the working sample was then taken into a metallic crucible and compacted carefully to reduce the rate of combustion. The combustion was carried out in a ballistic bomb calorimeter with appropriate corrections based on the standardization of the instrument with thermo-chemical

grade benzoic acid as the standard. The gross energy (heat of combustion) was calculated as cal/g.

VI. Estimation of crude fibre

Crude fibre was determined as the fraction remaining after digestion with standard solution of sulphuric acid and sodium hydroxide under carefully controlled condition.

A weighed quantity (2 g) of dry fat-free sample was taken in a spoutless conical flask fitted with a reflect condenser. Boiling water was added to it followed by addition of 25 ml sulphuric acid (10%). The content was mixed, the volume raised to 200 ml and boiled for 30 minutes. Acid solution was removed and the filtrate collected. The residue was then washed at least three times with boiling water. To this, was added 25 ml sodium hydroxide (10%) solution and the content diluted to the mark. It was boiled for another 30 minute. At the end of the boiling, filtering and washing were repeated. The resulting residue was transferred to a porcelain crucible, dehydrated using ethanol and dried to a constant weight at 100°C. This was incinerated for 2 hours in

a muffle furnace at 650°C. The fibre content was calculated on the basis of difference in weight of the residue after drying and ignition, the value (on moisture free basis) was expressed as per cent by weight.

VII. Estimation of chitin

Dried, powdered and weighed (2 g) sample was taken in a centrifuge tube. Sodium hydroxide (10%) solution was added to it, and the tube kept in a thermostat maintained at 60°C for 24 hours. It was then centrifuged at 4000 rpm for 15 minutes. The supernatant soluble protein was discarded and the residue dried to a constant weight. The chitin content was calculated on the basis of difference between initial and final weights, and expressed as per cent by weight.

RESULTS

Data on proximate composition and gross energy content of the feedstuffs have been given in Tables I-VI. Of the feedstuffs of plant origin, soybean oil cake (Glycine max) contained the maximum (51%) crude protein content. This was followed by groundnut oil cake (Arachis hypogea), with 47% crude protein. The protein in other oil cakes ranged between 33 to 40%. Among the

husks, brans and dusts, lentil dust (Lens culinaris) showed the maximum (30.7%) crude protein, followed by wheat bran (Triticum aestivum) with 15.8%, and pigeon pea dust (Cajanus cajan) with 14% crude protein. The protein values in other plant products, ranged between 4-10%. Among the other products, blood meal and sardine fish meal showed higher values of crude protein, amounting to 74% and 53%, respectively. Dried, silk worm pupae (Bombyx mori) and mantis shrimp (Squilla sp.) meal contained low protein, 40.6% and 33%, respectively. With the exception of soybean oil cake, the different oil cakes examined contained crude fat in the range of 6-10%.

The lowest value of fat observed in soybean oil cake could be ascribed to the use of solvent extraction procedure followed by the oil mill. Rice polish (Oryza sativa) contained 11.6% fat, whereas barley dust (Hordeum vulgare) and Bengal gram husk (Cicer arietinum) showed low values, 0.6% and 0.5%, respectively. The crude fat content in other products was in the range of 1.5 to 4.9%. In the animal by-products, the crude fat was maximum (21.7%) in silk worm pupae, followed by sardine fish meal (11.5%). The fat level in mantis shrimp meal and blood meal was low (1.8%) and comparable.

The moisture content in various feedstuffs ranged from 2.6 to 11.6%. The total ash content was highest (14.4%) in rice bran, followed by mantis shrimp meal (12.2%). The lowest (3.6%) ash was noted in barley dust. In the remaining feedstuffs, the ash content ranged between 4.3 and 10.5%.

Crude fibre content among the feedstuffs of plant origin was found to be the highest (42.3%) in Bengal gram husk. This was followed by cottonseed oil cake (Gossypium sativum) and pigeon pea dust with 28.4% and 25.7% crude fibre, respectively. Sesame oil cake (Sesamum indicum) contained 15.4% crude fibre, which is comparable to chitin content of silk worm pupae (13.2%). In other feedstuffs, the crude fibre content ranged between 7 and 11%.

The gross energy content of various ingredient studied depicted a wide range of energy density with the minimum (2962 cal/g) in mantis shrimp meal and maximum (5670 cal/g) in dried silk worm pupae. Sardine fish meal and slaughter-house waste contained more than 5000 cal/g of gross energy. Rice bran, pigeon pea husk, black gram husk and tapioca flour showed less than 4000 cal/g of gross energy. The oil cakes analysed, showed average values, ranging from 4492 to 4850 cal/g

of gross energy. Wheat bran had a comparable energy value with the oil cakes. Barley husk and Bengal gram husk showed similar values (4037 cal/g and 4036 cal/g). Rice polish and barseem showed a close range of energy density, comparable to different oil cakes.

Tables(VII-XVIII)contained additional information on the plant origin feedstuffs required for the INFIC retrieval system.

DISCUSSION

Although the usefulness of several artificial feedstuffs of plant and animal origin have been indicated for the spawn, fry and fingerlings of Indian major carps (Chakarborty et al., 1973; Jhingran, 1983; Mohanty et al., 1987; and Chattopadhyay and Konar, 1987) and the catfish (Venkatesh et al., 1986 and Niamat, 1982; and Niamat and Jafri 1984 a,b,) dietary formulations for these and other Indian cultivable fishes generally included combinations of a few conventional feedstuffs, mainly for supplementary feeding. Despite the wide variety of animal and plant origin feedstuffs, that are available at cheaper cost and that could possibly be incorporated in fish rations, very few have been nutritionally evaluated for the

above purpose. The use of locally available less expensive feed ingredients for carp polyculture in India has been stressed earlier in FAO technical report (Chow, 1982).

Since protein is by far the most important and often the most expensive dietary nutrient component required for growth, newer and cheaper protein sources, particularly those which do not compete as feed for other animals, are being scanned for use in fish feeds. Most fish species, depending upon their physiological needs, are generally known to require 20-60% crude protein in their diet (Hastings, 1979). The gross nutrient analysis of various feedstuffs carried out during the present study reveals the nutritive richness of a number of feedstuffs in terms of protein and other nutrients, and indicates to the feasibility of incorporating these low-cost ingredients in practical diets of our traditionally cultured fish species. It also points to the possibility where one ingredient could be replaced by the other. The use of conventional feed ingredients and a new generation of non-conventional products (SCP, plant protein concentrates, animal food processing wastes, etc.) as protein sources in complete diets for fish has been attempted by several workers in the past on various fish species (Tacon and Jackson,

1985; and Chow and Woo, 1989). Several of these attempts were directed to replace the fish meal portion of the diet (Cho et al., 1974; Spinellie et al., 1979; Viola et al., 1982; and Hilton, 1983), a commodity becoming expensive year after year with supplies not too dependable either. The analysis of various locally available feedstuffs indicate that byproducts like blood meal, fish meal, soybean meal, groundnut oil cake, containing over 45% crude protein, could be incorporated in high protein fish feeds. On the other hand, feedstuffs such as cottonseed meal, mustard oil cake, rocket salad oil cake, sesame oil cake, lentil dust and silkworm pupae could be utilised in formulating average protein diets.

In India and many of the south-east asian countries, where paddy is extensively cultivated, byproducts of rice mill, like rice polish and rice bran, are conventionally used in various combinations with different oil cakes as supplementary feed for carps and other fishes. The use of rice bran and low quality trash fish has been recommended in different rations as feed for certain species of freshwater catfish (Dehadrai and Thakur, 1980).

In formulating practical diets with desired level of protein, care must also be taken to consider

optimum level of energy (NAS-NRC 1983), so that the fish can make full utilization of available protein for their growth purposes. Ingredients like silkworm pupae, slaughter-house offal, fish meal, algal meal, certain oil cakes and brans can conveniently be used for the maintenance of energy level.

The fibre component of different feedstuffs can be useful in obtaining a proper nutrient balance as well as adequate water stability in the feed. Ingredients like cottonseed oil cake, sesame oil cake, pigeon pea dust, Bengal gram husk, silkworm pupae, with higher values of fibre/chitin, can be utilized for the above purpose.

It has been emphasized that in order to utilize protein efficiently for growth, other sources of energy such as fat and carbohydrate must also be present in fish diet (Halver, 1976; and Wilson and Halver, 1986). Since most fish have the ability to use fat and carbohydrate, to some extent, to spare protein as an energy source (Ringrose, 1971; Lee and Putnam, 1973; Takeuchi et al., 1978 a,b,c; Pieper and Pfeffer, 1979; Bergot, 1979 a,b. Takeuchi et al., 1979 ; Shimeno et al., 1980; Bromley 1980; Yu and Sinnhuber, 1981; Gatlin and Stickney, 1982, NAS-NRC, 1983; Stickney, 1984; and Strange, 1984), the inclusion of these nutrients are considered especially useful in deve-

loping feeds for young fish where protein needs for growth are the maximum (Hastings and Dupree, 1969; Mertz, 1972; Halver, 1976; and Wilson and Halver, 1986). As far as the requirement of fat for carp and catfish are concerned, it is reported to be 6-16% of the diet (Watanabe, 1982; and Stickney, 1984). Fats, being high energy nutrient, are known to be 85-90% digestible (Takeuchi et al., 1979; Austreng et al., 1980; NAS-NRC, 1981; Cho et al., 1985; Law, 1986; and Kirchgessner et al., 1986). Thus, besides fish meal, fat rich ingredients like mustard oil cake, rocket salad oil cake, sesame oil cake, rice polish and silkworm pupae could also be considered for inclusion in fish feeds at various levels.

The carbohydrate component of diet, besides providing energy, texturises the feed, acting as binder in case of expanded and compressed pellets. Ingredients like rice polish and tapioca flour appear to be of practical utility in this regard.

The result of the evaluation of gross energy content of locally available feedstuffs points to the nutritive richness of a good number of such ingredients. Some of the low-cost products like oil cakes, brans and husks, slaughter-house waste and silkworm pupae appeared exceedingly rich in calories.

The values obtained during the present study on the proximate composition and energy density of feedstuff were found generally comparable to those reported elsewhere (Ensminger and Olentine, 1980; and NAS-NRC; 1981, 1983).

It may, however, be pointed out that although many of the feedstuffs, scanned for their nutrient and energy content, could be useful in developing rations for cultivable fish species, the levels of inclusion of these ingredients must be determined keeping in view the nutritional requirements of the concerned species and also factors like the chemical characteristics, food value, digestibility, availability and overall cost of the feedstuffs.

SUMMARY

The survey of feedstuffs indicate that a great range of cheaper feedstuffs, both of plant and animal origin, are locally available. There being a greater variety of agricultural products that are cheaply and adequately available and their inclusion in fish diet could contribute to its cost-effectiveness. Analysis of nutritional qualities, based on proximate composition and calorific value, of these

locally available feedstuffs points to their nutritive richness. Several of the ingredients analysed were rich in their protein and gross energy contents. Some of the less expensive ingredients rich in protein could partly supplement the expensive or conventional protein sources in formulated feeds. Practical and economical rations can thus be formulated for cultivable fish species making greater use of locally available feedstuffs.




Table I : Proximate composition of feedstuffs (g/100g, as fed basis)

	Moisture	Crude Protein	Crude Fat	Ash	Crude Fibre
<u>Oil cakes</u>					
1. Mustard oil cake (<u>Brassica nigra</u>)	¹ 6.70±0.08 ² 6.50-7.05 ³ (6)	35.91 ± 1.00 31.44--39.10 (8)	9.63 ± 0.79 6.95 - 12.56 (6)	8.25 ± 0.09 8.05 - 8.65 (7)	10.45 ± 0.22 9.70 -10.94 (5)
2. Mustard oil cake (<u>Brassica campestris</u>)	2.64±0.14 2.35-3.20 (6)	37.94 ± 1.17 33.70 - 42.46 (8)	9.82 ± 0.12 9.30 - 10.09 (6)	7.79 ± 0.04 7.70 - 7.95 (6)	9.26 ± 0.20 8.50 - 9.70 (5)
3. Cotton seed oil cake (<u>Gossypium sativum</u>)	8.06±0.08 7.85-8.40 (6)	32.88 ± 1.27 29.56 - 37.50 (6)	5.65 ± 0.68 3.84 - 7.10 (5)	4.82 ± 0.21 4.45 - 5.50 (5)	28.35 ± 0.31 27.80 - 28.92 (4)
4. Rocket salad oil cake (<u>Eruca sativa</u>)	5.89±0.35 4.64-6.85 (6)	40.61 ± 1.46 35.60 - 46.75 (8)	7.98 ± 0.42 7.04 - 9.55 (6)	8.01 ± 0.19 7.50 - 8.50 (6)	9.63 ± 0.28 9.21 - 10.45 (4)
5. Soybean oil cake (<u>solvent</u> extd) (<u>Glycine max</u>)	7.05±0.46 5.80-8.55 (6)	51.05 ± 2.09 45.64 - 62.24 (8)	0.67 ± 0.16 0.20 - 1.36 (8)	7.13 ± 0.12 6.70 - 7.60 (6)	8.52 ± 0.17 8.20 - 9.10 (5)
6. Groundnut oil cake (<u>Arachis hypogaea</u>)	6.34±0.15 5.80-6.90 (6)	47.41 ± 1.15 43.42 -53.26 (8)	5.76 ± 0.95 3.14 - 8.05 (6)	6.75 ± 0.43 5.60 - 7.65 (4)	8.29 ± 0.27 7.75 - 9.04 (4)
7. Sesame oil cake (<u>Sesamum indicum</u>)	2.88±0.05 2.73-3.10 (6)	34.22 ± 0.97 29.02 - 37.54 (8)	6.72 ± 0.21 6.15 - 7.45 (6)	10.50 ± 0.14 10.15 - 11.00 (5)	15.43 ± 0.25 14.70 - 16.20 (5)
¹ ± SEM	² Range ; ³ Number of samples analysed.				

Table II : Proximate composition of feedstuffs (g/100 g, as fed basis)

Brans/Husk/Dust	Moisture	Crude Protein	Crude Fat	Ash	Crude Fibre
1. Pigeon pea dust (<u>Cajanus cajan</u>)	¹ 8.20 ± 0.41 ² 7.05 - 9.15 3(6)	13.96 ± 0.43 12.09 - 15.66 (8)	2.43 ± 0.37 0.70 - 3.49 (7)	5.43 ± 0.05 5.25 - 5.60 (6)	25.72 ± 0.28 25.16 - 26.50 (4)
2. Bengal gram husk (<u>Cicer arietinum</u>)	11.60 ± 0.26 10.50 - 12.15 (6)	4.28 ± 0.59 3.10 - 5.72 (4)	0.48 ± 0.15 0.18 - 0.65 (3)	4.28 ± 0.06 4.15 - 4.50 (5)	42.29 ± 0.21 41.70 - 42.69 (4)
3. Bengal gram dust (<u>Cicer arietinum</u>)	9.70 ± 0.45 8.40 - 11.00 (6)	9.58 ± 0.70 7.11 - 11.72 (8)	2.48 ± 0.21 1.70 - 3.10 (6)	4.26 ± 0.10 4.05 - 4.60 (5)	-
4. Lentil dust (<u>lens culinaris</u>)	8.02 ± 0.34 6.68 - 8.80 (6)	30.69 ± 0.92 27.06 - 34.26 (8)	1.49 ± 0.26 0.61 - 2.01 (5)	5.41 ± 0.07 5.15 - 5.65 (6)	7.11 ± 0.15 6.80 - 7.50 (5)
5. Barley dust (<u>Hordeum vulgare</u>)	10.11 ± 0.09 9.80 - 10.50 (6)	7.91 ± 0.22 7.20 - 8.96 (8)	0.61 ± 0.20 0.29 - 1.34 (5)	3.57 ± 0.13 3.15 - 4.00 (5)	8.60 ± 0.43 7.20 - 9.57 (5)
6. Wheat bran (<u>Triticum aestivum</u>)	10.86 ± 0.71 8.05 - 12.55 (6)	15.82 ± 0.41 14.20 - 17.33 (8)	4.89 ± 0.10 4.65 - 5.15 (5)	6.19 ± 0.08 5.95 - 6.50 (6)	10.26 ± 0.20 9.52 - 10.70 (5)
7. Rice polish (<u>Oryza sativa</u>)	9.06 ± 0.14 8.70 - 9.50 (5)	9.90 ± 0.49 8.25 - 11.30 (5)	11.57 ± 0.52 10.20 - 12.67 (4)	5.18 ± 0.17 4.90 - 5.90 (5)	-
8. Rice bran (<u>Oryza sativa</u>)	3.26 ± 0.17 2.80 - 3.80 (5)	5.87 ± 0.46 4.90 - 7.50 (5)	1.84 ± 0.12 1.57 - 2.10 (4)	14.40 ± 0.35 13.50 - 15.00 (4)	-

¹ ± SEM ; ² Range ; ³ Number of samples analysed

Table III : Proximate composition of feedstuffs (g/100 g, as fed basis)

		Moisture	Crude Protein	Crude Fat	Ash	Crude Fibre
<u>Miscellaneous</u>						
1. Sardine fish meal (<u>Sardinella</u> sp.)	¹	6.52 + 0.30	53.07 + 1.32	11.48 + 0.31	7.08 + 0.14	
	²	5.60 - 7.20	48.93 - 58.39	10.50 - 12.30	6.60 - 7.35	-
	³	(5)	(7)	(5)	(5)	
2. Blood meal		7.82 + 0.32	74.37 + 1.47	1.76 + 0.17	6.42 + 0.14	
		6.80 - 8.70	70.00 - 78.90	1.30 - 2.10	6.00 - 6.85	-
		(5)	(6)	(4)	(5)	
3. Mantis shrimp meal (<u>Squilla</u> sp)		5.20 + 0.29	33.18 + 1.82	1.81 + 0.19	12.21 + 0.19	
		4.70 - 5.70	29.90 - 38.20	1.30 - 2.20	11.85 - 12.50	-
4. Silkworm Pupae (<u>Bombyx mori</u>)		2.52 + 0.18	40.69 + 1.24	21.73 + 0.29	4.20 + 0.12	13.23 + 0.20
		2.21 - 2.63	38.11 - 41.01	20.89 - 22.33	3.82 - 4.78	12.32 - 14.29
¹ + SEM ; ² Range ; ³ Number of samples analysed						

Table IV : Gross Energy content of the feedstuffs

Feedstuffs	Gross Energy (cal/g)
<u>Oil Cakes</u>	
1. Mustard oil cake (<u>Brassica nigra</u>)	4748.5 \pm 69.51 (4501.0 - 4894.0)
2. Mustard oil cake (<u>Brassica campestris</u>)	4850.2 \pm 27.67 (4763.0 - 4938.0)
3. Cottonseed oil cake (<u>Gossypium sativum</u>)	4719.0 \pm 19.67 (4675.0 - 4763.0)
4. Rocket salad oil cake (<u>Eruca sativa</u>)	4492.0 \pm 37.66 (4413.0 - 4632.0)
5. Soybean oil cake (<u>Glycine max</u>)	4559.4 \pm 8.48 (4544.0 - 4588.0)
6. Groundnut oil cake (<u>Arachis hypogea</u>)	4657.8 \pm 22.22 (4588.0 - 4719.0)
7. Sesame oil cake (<u>Sesamum indicum</u>)	4596.8 \pm 59.39 (4544.0 - 4807.0)

+ SEM of 5 samples

Values in parentheses indicate the range

Table V : Gross Energy content of the feedstuffs

Feedstuffs	Gross Energy (cal/g)
<u>Brans/Husk/Dust</u>	
1. Rice bran (<u>Oryza sativa</u>)	3502.2 \pm 21.69 (3452.0 - 3583.0)
2. Wheat bran (<u>Triticum aestivum</u>)	4518.2 \pm 35.55 (4413.0 - 4632.0)
3. Pigeon pea husk (<u>Cajanus cajan</u>)	3800.5 \pm 24.22 (3730. - 3899.0)
4. Lentil husk (<u>Lens culinaris</u>)	4360.8 \pm 25.45 (4282.0 - 4413.0)
5. Lentil dust (<u>Lens culinaris</u>)	4334.8 \pm 25.65 (4238.0 - 4370.0)
6. Pigeon pea dust (<u>Cajanus cajan</u>)	4299.4 \pm 50.94 (4151.0 - 4413.0)
7. Barley husk (<u>Hordeum vulgare</u>)	4036.0 \pm 29.75 (3933.0 - 4107.0)
8. Bengal gram husk (<u>Cicer arietinum</u>)	4037.4 \pm 22.27 (3976.0 - 4107.0)
9. Black gram husk (<u>Phaseolus mungo</u>)	3810.2 \pm 25.45 (3758.0 - 3889.0)

\pm SEM of 5 samples

Values in parentheses indicate the range

Table VI : Gross Energy content of the feedstuffs

Feedstuffs	Gross Energy (cal/g)
<u>Miscellaneous</u>	
1. sardine fish meal (<u>Sardinella</u> sp.)	5216.2 \pm 55.93 (5037.0 - 5377.0)
2. Mantis shrimp meal (<u>Squilla</u> sp.)	2962.4 \pm 21.42 (2884.0 - 3015.0)
3. Slaughterhouse waste(offal)	5366.3 \pm 85.86 (5025.5 - 5462.5)
4. Silkworm pupae <u>Bombyx mori</u>	5670.0 \pm 39.78 5381.2 - 5810.7
5. Algal meal (<u>Spirulina</u> sp.)	4850.7 \pm 110.85 (4544.8 - 5069.2)
6. Tapioca flour (<u>Manihot utilissima</u>)	3915.5 \pm 32.78 (3845.6 - 4020.4)
7. Rice polish (<u>Oryza sativa</u>)	4579.6 \pm 15.34 (4550.0 - 4630.0)
9. Barseem (<u>Trifolium alexandrium</u>)	4559.36 \pm 29.24

\pm SEM of 5 samples

values in parentheses indicate the range

Table -VII

International Source Form for Composition of Feeds

Project No. 645 Experiment No FSA-1 Leader Prof.A.K.Jafri

(Please print)

Laboratory name Fish Nutrition Research Laboratory

Address Department of Zoology,
Aligarh Muslim University, Aligarh, India

Laboratory director

Laboratory sample No. 1

Origin and Name of Sample

Date originally collected Month March Day Monday Year 1987

Country India

Region Northern India

State, province or department

County or district Aligarh

Literature reference ICAR-Handbook of Agriculture

Food Information Center
Food and Agriculture Organization
of the United Nations (FAO, AGA)
Via delle Terme di Caracalla
00100 Rome, ItalyMail
duplicate to

Read instructions before filling in form.

very thin	thin	thrifty	fat	very fat
Percent of test ingredient in ration fed (100.0% dry matter)				
Ad Libitum feeding	✓	Controlled feeding		
Feed fed alone	✓	Feed not fed alone, digestion by difference		
Method: Total feces collection		Feces indicator	✓	

Physical form of feed Moist cake.

Basis for calculating level of feeding:	NE	ME	DE	TDN
Length of trial: Preliminary days	03	Collection days	03	
Daily dry matter consumed	g	kg		
Daily dry matter offered	g	kg		
Dailyorts or weighback	g	kg		
Temperature: Degrees C (indicate if negative)	0	26		
Pressure: millimeters				
Humidity: percent				
Card 40				

Dry Matter

Dry matter of sample on an as fed basis

Dry Matter	Method of analysis, if any, by another method but under same conditions	Anal. code factor	Digestion coeff. %
101	9 2.9		

Dry Matter Basis on Which Analytical Data are Reported on This Form

NOTE: All analytical data on this sheet, except dry matter, must be expressed as indicated: a. as fed, partially dry, dry (100.0% dry matter). Where analytical data for one feed is partly on one basis and partly on another, use separate sheet for each basis or convert to same basis.

Check Basis one only	Fill in one only
As fed	102
Partially dry	103
Dry (100.0% dry matter)	104

Proximate Principles

Ash	0 7.1	Assoc. Official Agr. Chem 1960	Anal. code factor	Digestion coeff. %
Crude fiber	0 8.5	Wierde		
Ether extract	0 0.6	Assoc. Official Agr. Chem 1960		
Nitrogen-free extract	2 5.5	By difference		
Protein	5 1.0	Kjeldahl		
Nitrogen	6 2.5	do		
Nitrogen factor		Write in factor to convert to protein		

Energy

Gross energy	4 5.59	Bomb calorimeter	
Digestible	0 0.79	Harris Natl. Acad. Sci. Natl. Res. Council pub. 1411, 1966	
Metabolizable		do	
N-equilibrium metabolizable		do	
NE _m		Lalgreen J. Animal Sci. 27:793 1968	
NE gain		do	
NE lactating cows		Moe J. Dairy Sci. 52:928	

TON

Biological

Card 10

12				
17				
22				

Do not write in shaded areas.

For Code

24				
----	--	--	--	--

Date originally collected

Month March Day Monday Year 1987

Country India

Region Northern India

State, province or department

County or district Aligarh

Literature reference ICAR-Handbook of Agriculture

Class	Dry forage (cut and cured)	Pasture (including range plants)	Cut and fed green	Silage
Scientific name: Genus	Glycine			
Species	max			
Variety or kind	Type - Verde, Bansal,			
Common name of feed	Geri kalai			

International feed reference No. 58

Part of plant, animal or feed product Beans

Process undergone before fed to animal

Stage of maturity	75 - 110 days
Cutting or crop	1
Height when cut (centimeters)	0 180
Official grade or No.	16
Fertilizer: kg per hectare	N - - - - - P - - - - - K - - - - -
Days between last application and harvest	28

Purity %

Chief contaminant

Soil classification Alluvium

Surface texture Sandy

Digestibility Trial

Animal: Kind Fish (Labeo rohita)

Breed

Sex

Age: Years 0 + Months Weeks

Number of animals in treatment 10

Total number of determinations per feed per animal 2

Average weight of animals, kg

Physiological state: non-pregnant pregnant 1st 2/3 pregnant last 1/3

losing wt. maintaining wt. gaining wt. fattening

Card 30	12			
15				
18				

21				
19				

28				
26				
29				

30				
36				
37				

39				
40				
42				
43				
44				

45				
48				
49				
51				
53				
59				
65				
71				
74				
78				

Table - VIII

International Source Form for Composition of Feeds

Project No. **645** Experiment No **FSA-1** Leader **Prof. A.K.Jafri**

Laboratory name **Fish Nutrition Research Laboratory**

Address **Department of Zoology,
Aligarh Muslim University, Aligarh, India**

Laboratory director

Laboratory sample No. **2** For Code

Origin and Name of Sample

Date originally collected **March Monday Year 1987 30**

Country **India**

Region **Northern India** Altitude (meters) **184** 39 42 36

State, province or department **U.P.**

County or district **Aligarh**

Literature reference **ICAR-Handbook of Agriculture 51** 46 48

Class **Dry forage (cut and cured)** Pasture (including range plants) **Gossypium** Silage

Scientific name: Genus **Gossypium**

Species **herbaceum**

Variety or kind **320F, 216F**

Common name of feed **Kapas**

Part of plant, animal or feed product **Seeds (oil cake)**

Process undergone before fed to animal

Stage of maturity **240 days**

Culturing or crop **1** Height when cut (centimeters) **0100** 12 24

Official grade or No. **16** 20

Fertilizer: kg per hectare **N 0015 P 0012 K 0012**

Purity % **28**

Soil classification **Alluvium** 31

Surface texture **Sandy** 34

Digestibility Trial

Animal: Kind **Fish (Labeo rohita)**

Breed **Sex**

Age: Years **0** + Months **10** Weeks

Number of animals in treatment **10**

Total number of determinations per feed per animal **2** 29 30

Average weight of animals, kg **24.7**

Physiological state: non-pregnant **pregnant 1st 2/3** 36 37

losing wt **maintaining wt** **gaining wt** **fattening**

very thin ☐ thin ☐ thrifty ☐ fat ☐ very fat ☐

Percent of test ingredient in ration fed (100.0% dry matter)

Ad Libitum feeding ☒ Controlled feeding ☐

Feed fed alone ☒ Feed not fed alone, digestion by difference ☐

Method: Total feces collection ☐ feces indicator ☒

Physical form of feed **Moist cake.**

Basis for calculating level of feeding: NE ME DE TDN

Length of trial: Preliminary days **03** Collection days **03**

Daily dry matter consumed **g** **kg**

Daily dry matter offered **g** **kg**

Dailyorts or weightback **g** **kg**

Temperature: Degrees C (indicate if negative) **026**

Pressure: millimeters

Humidity: percent

Card 40

Dry Matter

Dry Matter: Method of analysis, if area is done by another method but under same basis

Digestion coeff. %

Anal. code factor

Dry Matter Basis on Which Analytical Data are Reported on This Form

NOTE: All analytical data on this sheet, except dry matter, must be expressed as indicated: i.e. on fed, partially dry, dry (100.0% dry matter). Where analytical data for one feed is partly on one basis and partly on another, use separate sheet for each basis or convert to same basis.

Proximate Principles

Check Basis one only ☒ At fed ☐ Partially dry ☐ Dry (100.0% dry matter) ☐

Fill in one only

Ash **048** 105

Crude fiber **283** 106

Ether extract **056** 107

Nitrogen-free extract **202** 108

Protein **328** 109

Nitrogen **625** 212

Nitrogen factor **625** 213

Energy **4719** kcal per kg

Crack energy **4719** 421

Digestible **4719** 422

Metabolizable **4719** 423

N-equilibrium **4719** 424

NE gain **4719** 426

NE lactating cows **4719** 427

TDN **4719** 429

Method of analysis, if analysis was done by another method but under other analyses

Digestion coeff. %

Anal. code factor

39 40 42 43 44

45 48 49 51 53 59 65 71 74 78

Read instructions before filling in form.

Feed Information Center
Food and Agriculture Organization
of the United Nations (FAO, AGA)
Viale delle Terme di Caracalla
00100 Rome, Italy

Table -IX

International Source Form for Composition of Feeds

Project No. 645 Experiment No. FSA-1 Investigator Prof. A.K. Jafri

Laboratory name Fish Nutrition Research Laboratory

Address Department of Zoology,
Aligarh Muslim University, Aligarh, India.

Laboratory director

Laboratory sample No. 3

Origin and Name of Sample
Date originally collected Month March Day Monday Year 1987
Country India
Region Northern India Altitude (meters) 184
State, province or department U.P.
County or district Aligarh
Literature reference ICAR-Handbook of Agriculture 51

Class Dry forage (cut and cured) Pasture (including range plants) Cut and fed green Silage

Scientific name: Genus Arachis
Species hypogaea
Variety or kind Punjab 1
Common name of feed Mangfuli, Cheena badam.

Part of plant, animal or feed product Pod (oil cake)

Process undergone before fed to animal Drying, grinding, sieving

Stage of maturity 95 days

Cutting or crop 1 Height when cut (centimeters) 0 0 2 5

Official grade or No. 16 N 00 1 5 P 00 3 0 K 00 2 5

Fertilizer: kg per hectare 20

Days between last application and harvest

Purity % 100 Chief contaminant

Soil classification Alluvium

Surface texture Loam/Sandy-loam

Digestibility Trial

Animal: Kind Fish (Labeo rohita)

Breed Sex

Age: Years 0 + Months 10 Weeks

Number of animals in treatment 10

Total number of determinations per feed per animal 2 period No.

Average weight of animals, kg or g 24.7

Physiological state: non-pregnant pregnant 1st 2/3 pregnant last 1/3

losing wt maintaining wt gaining wt fattening

Card 10

Do not write in shaded areas.

For Code

Card 20

Card 30

Card 40

Card 50

Card 60

Card 70

Card 80

Card 90

Card 100

Card 110

Card 120

Card 130

Card 140

Card 150

Card 160

Card 170

Card 180

Card 190

Card 200

Card 210

Card 220

Card 230

Card 240

Card 250

Card 260

Card 270

Card 280

Card 290

Card 300

Card 310

Card 320

Card 330

Card 340

Card 350

Card 360

Card 370

Card 380

Card 390

Card 400

Card 410

Card 420

Card 430

Card 440

Card 450

Card 460

Card 470

Card 480

Card 490

Card 500

Card 510

Card 520

Card 530

Card 540

Card 550

Card 560

Card 570

Card 580

Card 590

Card 600

Card 610

Card 620

Card 630

Card 640

Card 650

Card 660

Card 670

Card 680

Card 690

Card 700

Card 710

Card 720

Card 730

Card 740

Card 750

Card 760

Card 770

Card 780

Card 790

Card 800

Card 810

Card 820

Card 830

Card 840

Card 850

Card 860

Card 870

Card 880

Card 890

Card 900

Card 910

Card 920

Card 930

Card 940

Card 950

Card 960

Card 970

Card 980

Card 990

Card 1000

Card 1010

Card 1020

Card 1030

Card 1040

Card 1050

Card 1060

Card 1070

Card 1080

Card 1090

Card 1100

Card 1110

Card 1120

Card 1130

Card 1140

Card 1150

Card 1160

Card 1170

Card 1180

Card 1190

Card 1200

Card 1210

Card 1220

Card 1230

Card 1240

Card 1250

Card 1260

Card 1270

Card 1280

Card 1290

Card 1300

Card 1310

Card 1320

Card 1330

Card 1340

Card 1350

Card 1360

Card 1370

Card 1380

Card 1390

Card 1400

Card 1410

Card 1420

Card 1430

Card 1440

Card 1450

Card 1460

Card 1470

Card 1480

Card 1490

Card 1500

Card 1510

Card 1520

Card 1530

Card 1540

Card 1550

Card 1560

Card 1570

Card 1580

Card 1590

Card 1600

Card 1610

Card 1620

Card 1630

Card 1640

Card 1650

Card 1660

Card 1670

Card 1680

Card 1690

Card 1700

Card 1710

Card 1720

Card 1730

Card 1740

Card 1750

Card 1760

Card 1770

Card 1780

Card 1790

Card 1800

Card 1810

Card 1820

Card 1830

Card 1840

Card 1850

Card 1860

Card 1870

Card 1880

Card 1890

Card 1900

Card 1910

Card 1920

Card 1930

Card 1940

Card 1950

Card 1960

Card 1970

Card 1980

Card 1990

Card 2000

Card 2010

Card 2020

Card 2030

Card 2040

Card 2050

Card 2060

Card 2070

Card 2080

Card 2090

Card 2100

Card 2110

Card 2120

Card 2130

Card 2140

Card 2150

Card 2160

Card 2170

Card 2180

Card 2190

Card 2200

Card 2210

Card 2220

Card 2230

Card 2240

Card 2250

Card 2260

Card 2270

Card 2280

Card 2290

Card 2300

Card 2310

Card 2320

Card 2330

Card 2340

Card 2350

Card 2360

Card 2370

Card 2380

Card 2390

Card 2400

Card 2410

Card 2420

Card 2430

Card 2440

Card 2450

Card 2460

Card 2470

Card 2480

Card 2490

Card 2500

Card 2510

Card 2520

Card 2530

Card 2540

Card 2550

Card 2560

Card 2570

Card 2580

Card 2590

Card 2600

Card 2610

Card 2620

Card 2630

Card 2640

Card 2650

Card 2660

Card 2670

Card 2680

Card 2690

Card 2700

Card 2710

Card 2720

Card 2730

Card 2740

Card 2750

Card 2760

Card 2770

Card 2780

Card 2790

Card 2800

Card 2810

Card 2820

Card 2830

Card 2840

Card 2850

Card 2860

Card 2870

Card 2880

Card 2890

Card 2900

Card 2910

Card 2920

Card 2930

Card 2940

Card 2950

Card 2960

Card 2970

Card 2980

Card 2990

Card 3000

Card 3010

Card 3020

Card 3030

Card 3040

Card 3050

Card 3060

Card 3070

Card 3080

Card 3090

Card 3100

Card 3110

Card 3120

Card 3130

Card 3140

Card 3150

Card 3160

Card 3170

Card 3180

Card 3190

Card 3200

Card 3210

Card 3220

Card 3230

Card 3240

Card 3250

Card 3260

Card 3270

Card 3280

Card 3290

Card 3300

Card 3310

Card 3320

Card 3330

Card 3340

Card 3350

Card 3360

Card 3370

Card 3380

Card 3390

Card 3400

Card 3410

Card 3420

Card 3430

Card 3440

Card 3450

Card 3460

Card 3470

Card 3480

Card 3490

Card 3500

Card 3510

Card 3520

Card 3530

Card 3540

Card 3550

Card 3560

Card 3570

Card 3580

Card 3590

Card 3600

Card 3610

Card 3620

Card 3630

Card 3640

Card 3650

Card 3660

Card 3670

Card 3680

Card 3690

Card 3700

Card 3710

Card 3720

Card 3730

Card 3740

Card 3750

Card 3760

Card 3770

Card 3780

Card 3790

Card 3800

Card 3810

Card 3820

Card 3830

Card 3840

Card 3850

Card 3860

Card 3870

Card 3880

Card 3890

Card 3900

Card 3910

Card 3920

Card 3930

Card 3940

Card 3950

Card 3960

Card 3970

Card 3980

Card 3990

Card 4000

Card 4010

Card 4020

Card 4030

Card 4040

Card 4050

Card 4060

Card 4070

Card 4080

Card 4090

Card 4100

Card 4110

Card 4120

Card 4130

Card 4140

Card 4150

Card 4160

Card 4170

Card 4180

Card 4190

Card 4200

Card 4210

Card 4220

Card 4230

Card 4240

Card 4250

Card 4260

Card 4270

Card 4280

Card 4290

Card 4300

Card 4310

Card 4320

Card 4330

Card 4340

Card 4350

Card 4360

Card 4370

Card 4380

Card 4390

Card 4400

Card 4410

Card 4420

Card 4430

Card 4440

Card 4450

Card 4460

Card 4470

Card 4480

Card 4490

Card 4500

Card 4510

Card 4520

Card 4530

Card 4540

Card 4550

Card 4560

Card 4570

Card 4580

Card 4590

Card 4600

Card 4610

Card 4620

Card 4630

Card 4640

Card 4650

Card 4660

Card 4670

Card 4680

Card 4690

Card 4700

Card 4710

Card 4720

Card 4730

Card 4740

Card 4750

Card 4760

Card 4770

Card 4780

Card 4790

Card 4800

Card 4810

Card 4820

Card 4830

Card 4840

Card 4850

Card 4860

Card 4870

Card 4880

Card 4890

Card 4900

Card 4910

Card 4920

Card 4930

Card 4940

Card 4950

Card 4960

Card 4970

Card 4980

Card 4990

Card 5000

Card 5010

Card 5020

Card 5030

Card 5040

Card 5050

Card 5060

Card 5070

Card 5080

Card 5090

Card 5100

Card 5110

Card 5120

Card 5130

Card 5140

Card 5150

Card 5160

Card 5170

Card 5180

Card 5190

Card 5200

Card 5210

Card 5220

Card 5230

Card 5240

Card 5250

Card 5260

Card 5270

Card 5280

Card 5290

Card 5300

Card 5310

Card 5320

Card 5330

Card 5340

Card 5350

Card 5360

Card 5370

Card 5380

Card 5390

Card 5400

Card 5410

Card 5420

Card 5430

Card 5440

Card 5450

Card 5460

Card 5470

Card 5480

Card 5490

Card 5500

Card 5510

Card 5520

Card 5530

Card 5540

Card 5550

Card 5560

Card 5570

Card 5580

Card 5590

Card 5600

Card 5610

Card 5620

Card 5630

Card 5640

Card 5650

Card 5660

Card 5670

Card 5680

Card 5690

Card 5700

Card 5710

Card 5720

Card 5730

Card 5740

Card 5750

Card 5760

Card 5770

Card 5780

Card 5790

Card 5800

Card 5810

Card 5820

Card 5830

Card 5840

Card 5850

Card 5860

Card 5870

Card 5880

Card 5890

Card 5900

Card 5910

Card 5920

Card 5930

Card 5940

Card 5950

Card 5960

Card 5970

Card 5980

Card 5990

Card 6000

Card 6010

Card 6020

Card 6030

Card 6040

Card 6050

Card 6060

Card 6070

Card 6080

Card 6090

Card 6100

Card 6110

Card 6120

Card 6130

Card 6140

Card 6150

Card 6160

Card 6170

Card 6180

Card 6190

Card 6200

Card 6210

Card 6220

Card 6230

Card 6240

Card 6250

Card 6260

Card 6270

Card 6280

Card 6290

Card 6300

Card 6310

Card 6320

Card 6330

Card 6340

Card 6350

Card 6360

Card 6370

Card 6380

Card 6390

Card 6400

Card 6410

Card 6420

Card 6430

Card 6440

Card 6450

Card 6460

Card 6470

Card 6480

Card 6490

Card 6500

Card 6510

Card 6520

Card 6530

Card 6540

Card 6550

Card 6560

Card 6570

Card 6580

Card 6590

Card 6600

Card 6610

Card 6620

Card 6630

Card 6640

Card 6650

Card 6660

Card 6670

Card 6680

Card 6690

Card 6700

Card 6710

Card 6720

Card 6730

Card 6740

Card 6750

Card 6760

Card 6770

Card 6780

Card 6790

Card 6800

Card 6810

Card 6820

Card 6830

Card 6840

Card 6850

Card 6860

Card 6870

Card 6880

Card 6890

Card 6900

Card 6910

Card 6920

Card 6930

Card 6940

Card 6950

Card 6960

Card 6970

Card 6980

Card 6990

Card 7000

Card 7010

Card 7020

Card 7030

Card 7040

Card 7050

Card 7060

Card 7070

Card 7080

Card 7090

Card 7100

Card 7110

Card 7120

Card 7130

Card 7140

Card 7150

Card 7160

Card 7170

Card 7180

Card 7190

Card 7200

Card 7210

Card 7220

Card 7230

Card 7240

Card 7250

Card 7260

Card 7270

Card 7280

Card 7290

Card 7300

Card 7310

Card 7320

Card 7330

Card 7340

Card 7350

Card 7360

Card 7370

Card 7380

Card 7390

Card 7400

Card 7410

Card 7420

Card 7430

Card 7440

Card 7450

Card 7460

Card 7470

Card 7480

Card 7490

Card 7500

Card 7510

Card 7520

Card 7530

Card 7540

Card 7550

Card 7560

Card 7570

Card 7580

Card 7590

Card 7600

Card 7610

Card 7620

Card 7630

Card 7640

Card 7650

Card 7660

Card 7670

Card 7680

Card 7690

Card 7700

Card 7710

Card 7720

Card 7730

Card 7740

Card 7750

Card 7760

Card 7770

Card 7780

Card 7790

Card 7800

Card 7810

Card 7820

Card 7830

Card 7840

Card 7850

Card 7860

Card 7870

Card 7880

Card 7890

Card 7900

Card 7910

Card 7920

Card 7930

Card 7940

Card 7950

Card 7960

Card 7970

Card 7980

Card 7990

Card 8000

Card 8010

Card 8020

Card 8030

Card 8040

Card 8050

Card 8060

Card 8070

Card 8080

Card 8090

Card 8100

Card 8110

Card 8120

Card 8130

Card 8140

Card 8150

Card 8160

Card 8170

Card 8180

Card 8190

Card 8200

Card 8210

Card 8220

Card 8230

Card 8240

Card 8250

Card 8260

Card 8270

Card 8280

Card 8290

Card 8300

Card 8310

Card 8320

Card 8330

Card 8340

Card 8350

Card 8360

Card 8370

Card 8380

Card 8390

Card 8400

Card 8410

Card 8420

Card 8430

Card 8440

Card 8450

Card 8460

Card 8470

Card 8480

Card 8490

Card 8500

Card 8510

Card 8520

Card 8530

Card 8540

Card 8550

Card 8560

Card 8570

Card 8580

Card 8590

Card 8600

Card 8610

Card 8620

Card 8630

Card 8640

Card 8650

Card 8660

Card 8670

Card 8680

Card 8690

Card 8700

Card 8710

Card 8720

Card 8730

Card 8740

Card 8750

Card 8760

Card 8770

Card 8780

Card 8790

Card 8800

Card 8810

Card 8820

Card 8830

Card 8840

Card 8850

Card 8860

Card 8870

Card 8880

Card 8890

Card 8900

Card 8910

Card 8920

Card 8930

Card 8940

Card 8950

Card 8960

Card 8970

Card 8980

Card 8990

Card 9000

Card 9010

Card 9020

Card 9030

Card 9040

Card 9050

Card 9060

Card 9070

Card 9080

Card 9090

Card 9100

Card 9110

Card 9120

Card 9130

Card 9140

Card 9150

Card 9160

Card 9170

Card 9180

Card 9190

Card 9200

Card 9210

Card 9220

Card 9230

Card 9240

Card 9250

Card 9260

Card 9270

Card 9280

Card 9290

Card 9300

Card 9310

Card 9320

Card 9330

Card 9340

Card 9350

Card 9360

Card 9370

Card 9380

Card 9390

Card 9400

Card 9410

Card 9420

Card 9430

Card 9440

Card 9450

Card 9460

Card 9470

Card 9480

Card 9490

Card 9500

Card 9510

Card 9520

Card 9530

Card 9540

Card 9550

Card 9560

Card 9570

Card 9580

Card 9590

Card 9600

Card 9610

Card 9620

Card 9630

Card 9640

Card 9650

Card 9660

Card 9670

Card 9680

Card 9690

Table X

International Source Form for Composition of Feeds

(2 pages)

Project No. 645 Experiment No. FSA-1 Investigator Prof. A.K. Jafri

Laboratory name Fish Nutrition Research Laboratory

Address Department of Zoology,
Aligarh Muslim University, Aligarh, India

Laboratory director

Laboratory sample No. 4 For Code

Origin and Name of Sample

Date originally collected Month March Day Monday Year 1987

Country India

Region Northern India Altitude (meters) 184 39 42 36

State, province or department U.P. 46

County or district Aligarh 48

Literature reference ICAR-Handbook of Agriculture 51

Class Dry forage (cut and cured) Pasture (including range plants) Cut and fed green Silage

Scientific name: Genus Sesamum

Species indicum

Variety or kind T-85, N-32

Common name of feed T11

Part of plant, animal or feed product Seed (oil cake)

Process undergone before fed to animal

Stage of maturity 135 days

Cutting or crop 1 Height when cut (centimeters) 0 1 0 0

Official grade or No. 16 20 24 28

Fertilizer: kg per hectare N 0 0 2 5 P 0 0 2 0 K 0 0 0 8

Days between last application and harvest

Purity % Chief contaminant

Soil classification Field classification

Digestibility trial Surface texture Sandy loam-heavy black soil

Animal: Kind Fish (Labeo rohita)

Breed Sex

Age: Years 0 + Months Weeks

Number of animals in treatment 10

Total number of determinations per feed per animal 2 period No. 29

Average weight of animals, kg or g 2 4 7

Physiological state: non-pregnant pregnant 1st 2/3 3rd

Losing wt. maintaining wt. gaining wt. fattening

very thin	thin	thrifty	fat	very fat
Percent of test ingredient in ration fed (100.0% dry matter)				
Ad Libitum feeding	<input checked="" type="checkbox"/>	Controlled feeding		
Feed fed alone	<input checked="" type="checkbox"/>	Feed not fed alone, digestion by difference		
Method: Total feces collection		Feces indicator	<input checked="" type="checkbox"/>	

Physical form of feed Moist cake.

Basis for calculating level of feeding:				
Length of trial:	Preliminary days	NE	ME	TDN
Daily dry matter consumed	g	103	Collection days	0 3
Daily dry matter offered	g			
Dailyorts or weightback	g			
Temperature: Degrees C (indicate if negative)		0 2 6		
Pressure: millimeters				
Humidity: percent				

Dry Matter

Dry Matter	Method of analyses, if one method put under other analyses	Anal. code factor	Digestion coeff. %
97.1	Above 1050 C or in vacuo		

Dry Matter Basis on Which Analytical Data are Reported on This Form

NOTE: All analytical data on this sheet, except dry matter, must be expressed on indicated basis: as fed, control dry, dry (100.0% dry matter). Where analytical data for one feed is partly on one basis and partly on another, use separate sheet for each basis or convert to same basis.

Check Basis	Fill in one only
As fed	<input checked="" type="checkbox"/>
Partially dry	<input type="checkbox"/>
Dry (100.0% dry matter)	<input type="checkbox"/>

Proximate Principles

Ash	105	10.5	Assoc. Official Agr. Chem. 1960	Anal. code factor	Digestion coeff. %
Crude fiber	106	15.4	Weende		
Ether extract	107	06.7	Assoc. Official Agr. Chem. 1960		
Nitrogen-free extract	108	30.2	By difference		
Protein	109	34.2	Kjeldahl		
Nitrogen	212		do		
Nitrogen factor	213	6.25	Write in factor to convert to protein		

Energy

Gross energy	421	4596	Bomb calorimeter	Anal. code factor	Digestion coeff. %
Digestible	422		Hess, Natl. Acad. Sci. Natl. Res. Council pub. 141, 1960		
Metabolizable	423		do		
N-equilibrium metabolizable	424		do		
ME _m	426		Lofgreen J. Animal Sci. 27:793, 1968		
ME _g gain	427		do		
ME lactating cows	430		Moe J. Dairy Sci. 52:928		

TDN

TDN	429		Biological
-----	-----	--	------------

Table XI

International Source Form for Composition of Feeds

IP 2-10-57 (10/1)

Project No. 645 Experiment No. FSA-1 Leader Prof. A.K. Jafri

Laboratory name Fish Nutrition Research Laboratory

Address Department of Zoology,
Aligarh Muslim University, Aligarh, India

Laboratory director

Laboratory sample No.

5

Origin and Name of Sample

Date originally collected March Day Monday Year 1987

Country

Region Northern India

State, province or department U.P.

County or district Aligarh

Literature reference ICAR-Handbook of Agriculture

Class Dry forage (cut and cured)

Pasture (including range plants)

Cut and fed green

Silage

Scientific name: Genus Brassica

Species nigra

Variety or kind Type- 151

Common name of feed Sarson

Part of plant, animal or feed product Seed (oil cake)

Process undergone before fed to animal Drying, grinding, sieving

Stage of maturity 155 days

Cutting or crop 1

Height when cut (centimeters) 0175

Official grade or No. 16

Fertilizer: kg per hectare N 0 0 40 P - - - - K - - - -

Days between last application and harvest

Purity %

Soil classification Alluvium

Surface texture Light to heavy loam.

Digestibility Trial

Animal: Kind Fish (Labeo rohita)

Breed -

Sex -

Age: Years 0 + Months Weeks

Number of animals in treatment 10

Total number of determinations per feed per animal 2

Average weight of animals, kg 24.7

Physiological state: non-pregnant pregnant 1st 2/3 pregnant last 1/3

losing wt. maintaining wt. gaining wt. fattening

Card 10

12 22
17Do not write
in shaded
areas

For Code

24

36

42

46

48

51

58

64

70

76

82

88

94

100

106

112

118

124

130

136

142

148

154

160

166

172

178

184

190

196

202

208

214

220

226

232

238

very thin thin thrifty fat very fat
Percent of test ingredient in ration fed (100.0% dry matter)
Ad Libitum feeding ☒ Controlled feeding ☐
Feed fed alone ☒ Feed not fed alone, digestion by difference ☐
Method: Total feces collection ☐ feces indicator ☒

Physical form of feed Moist cake.

Basis for calculating level of feeding: NE ME DE TDN
Length of trial: Preliminary days 03 Collection days 03
Daily dry matter consumed g kg
Daily dry matter offered g kg
Daily orls or weighback g kg
Temperature: Degrees C (indicate if negative) 026
Pressure: millimeters
Humidity: percent
Card 40

Dry Matter

Dry Matter: ☒ Method of analysis, if one of the following: ☐ by another method put under other analyses
Dry matter of sample on a fed basis 101 93.3 Above 100% C or in vacuo

Dry Matter Basis on Which Analytical Data are Reported on This Form

NOTE: All analytical data on this sheet, except dry matter, must be expressed as indicated: i.e.: on fed, partial v dry, dry (100.0% dry matter). Where analytical data is one feed it partly on one basis and partly on another use separate sheet for each basis or convert to same basis
Check Basis one only
As fed ☒ 102
Partially dry ☐ 103
Dry (100.0% dry matter) ☐ 104

Proximate Principles

Ash 105 08.2
Crude fiber 106 10.4
Ether extract 107 09.6
Nitrogen-free extract 108 29.0
Protein 109 33.9
Nitrogen 212 do
Nitrogen factor 213 Write in factor to convert to protein
Energy kcal per kg
Gross energy 421 4748
Digestible 422 0086
Metabolizable 423 do
N-equilibrium metabolizable 424 do
NE_m 426 do
NE gain 427 do
NE lactating cows 430 do
TDN 429 doAnal. code factor
Digestion coeff %
Bomb calorimeter
Harris Natl Acad Sci Natl Res Council pub 1411, 1965
do
do
Lofgreen J Animal Sci 27:793, 1968
do
Moe J Dairy Sci 52:928

Biological

Table XII

International Source Form for Composition of Feeds

Project No. 645 Experiment No. FSA-1 Tester Prof. A.K. Jafri

Card 10

Laboratory name Fish Nutrition Research Laboratory

Address Department of Zoology,

Alligarh Muslim University, Alligarh, India

Laboratory director

Laboratory sample No. 6

Origin and Name of Sample

Date originally collected Month March Day Monday Year 1987

Country India

Region Northern India

State, province or department

County or district Alligarh

Literature reference ICAR-Handbook of Agriculture

Class Dry forage (cut and cured)

Scientific name: Genus Clitor

Variety or kind H 208, T 3, K 4

Common name of feed Chana, Chola

Part of plant, animal or feed product Seed (Husk, dust)

Process undergone before fed to animal Drying, grinding, sieving

Stage of maturity 150 days

Cutting or crop 1

Official grade or No. 16

Fertilizer: kg per hectare N 0015 P 0025 K 28

Purity % Days between last application and harvest

Soil classification Alluvium

Surface texture Clayey-loam

Digestibility Trial

Animal: Kind Fish (Labeo rohita)

Breed

Age: Years 0 + Months 10

Number of animals in treatment 10

Total number of determinations per feed per animal 2

Average weight of animals, kg 24.7

Physiological state: non-pregnant pregnant 1st 2/3 pregnant last 1/3

losing wt maintaining wt gaining wt fattening

Sex

Do not write in shaded areas.

For Code

Card 10

Card 10

Card 10

Card 10

Card 10

Card 10

Card 10

Card 10

Card 10

Card 10

very thin	thin	thrifty	fat	very fat
Percent of test ingredient in ration fed (100.0% dry matter)				
Ad Libitum feeding	<input checked="" type="checkbox"/>	Controlled feeding		
Feed fed alone	<input checked="" type="checkbox"/>	Feed not fed alone, digestion by difference		
Method: Total feces collection		Feces indicator	<input checked="" type="checkbox"/>	

Physical form of feed Moist cake.

Basis for calculating level of feeding:				
Length of trial:	Preliminary days	03	ME	TDN
Daily dry matter consumed	g		kg	
Daily dry matter offered	g		kg	
Dailyorts or weighback	g		kg	
Temperature: Degrees C (indicate if negative)	026			
Pressure: millimeters				
Humidity: percent				

Dry Matter

Dry matter of sample on as fed basis	101	88.4	4
Method of analysis, if any, by another method put under other analyses	Above 105°C or in vacuum		
Anal. code			
Digest. or coeff. factor			

Dry Matter Basis on Which Analytical Data are Reported on This Form

NOTE: All analytical data on this sheet, except dry matter, must be expressed as indicated: i.e. as fed, partially dry, dry (100.0% dry matter). Where analytical data for one feed is partly on one basis and partly on another use separate sheet for each basis or convert to same basis.

Proximate Principles

Ash	105	04.2	2
Crude fiber	106	42.2	2
Ether extract	107	0.4	2
Nitrogen-free extract	108	37.0	2
Protein	109	4.2	2
Nitrogen	212	6.2	5
Nitrogen factor	213	6.2	5

Energy

Gross energy	421	40.3	7
Digestible	422		
Metabolizable	423		
Non-equilibrium metabolizable	424		
NE _m	426		
NE gain	427		
NE lactating cows	430		

TDN

TDN	429		
-----	-----	--	--

Table XIII

International Source Form for Composition of Feeds

Project No. 645 Experiment No. PSA-1 Investigator Prof. A.K. Jafri

Laboratory name Fish Nutrition Research Laboratory

Address Department of Zoology,

Alligarh Muslim University, Alligarh, India

Laboratory director

Laboratory sample No. 7

Origin and Name of Sample

Date originally collected Month March Day Monday Year 1987

Country India

Region Northern India

State, province or department U.P.

County or district Alligarh

Literature reference ICAR-Handbook of Agriculture

Class Dry large (cut and cured)

Scientific name: Genus Lens

Species culinaris

Variety or kind T-36, Pant 209

Common name of feed Musur

Part of plant, animal or feed product Seed (Dust)

Process undergone before fed to animal Drying, grinding, sieving

Stage of maturity 100-120 days

Cutting or crop 1

Height when cut (centimeters) 00 3 0

Official grade or No. 16

Fertilizer: kg per hectare N

Days between last application and harvest

Purity %

Soil classification Alluvium

Digestibility Trial

Animal: Kind Fish (Labeo rohita)

Breed

Age: Years 0 + Months 10

Number of animals in treatment 10

Total number of determinations per feed per animal 2

Average weight of animals, kg

Physiological state: non-pregnant

losing wt

maintaining wt

gaining wt

fattening

Card 10

12

17

22

Do not write in shaded areas.

For Code

24

30

36

42

46

48

51

58

64

70

76

82

88

94

100

106

112

118

124

130

136

142

148

154

160

166

172

178

184

190

196

202

208

214

220

226

232

238

very thin	thin	thrifty	fat	very fat
Percent of test ingredient in ration fed (100.0% dry matter)				
Ad Libitum feeding	Controlled feeding			
Feed fed alone	Feed not fed alone, digestion by difference			
Method: Total feces collection	Feces indicator			

Physical form of feed Moist cake.

Basis for calculating level of feeding:	NE	ME	DE	TDN
Length of trial: Preliminary days	03			
Daily dry matter consumed	g	kg		
Daily dry matter offered	g	kg		
Dailyorts or weighback	g	kg		
Temperature: Degrees C (indicate if negative)				
Pressure: millimeters				
Humidity: percent				

Dry Matter	Dry Matter	Method of analysis, if any	Anal. code factor
on an as fed basis	on an as fed basis	by another method but under same conditions	
91.9	91.9	Above 105°C or in vacuum	

Dry Matter Basis on Which Analytical Data are Reported on This Form

Check one only	Fill in one only
Basis	As fed
Partially dry	102
Dry (100.0% dry matter)	103
	104

Proximate Principles

As	105	0.54	Asoc. Official Agr. Chem 1960
Crude fiber	106	0.71	Weende
Ether extract	107	0.14	Asoc. Official Agr. Chem 1960
Nitrogen-free extract	108	4.72	By difference
Protein	109	3.06	Kjeldahl
Nitrogen	212	6.25	do
Nitrogen factor	213	6.25	Write in factor to convert to protein

Energy

Gross energy	421	43.60	Bomb calorimeter
Digestible	422		Harris Natl. Acad. Sci. Natl. Res. Council pub 1411 1966
Metabolizable	423		do
Metabolizable	424		do
NE _m	426		Lalgreen J. Animal Sci. 27:793 1968
NE gain	427		do
NE lactating cows	430		Moe J. Dairy Sci. 52:928
TDN	429		Biological

Food Information Center
of the United Nations (FAO) AGAR
Via delle Terme di Caracalla
00100 Rome, Italy

Read instructions before filling in form.

Table XIV
International Source Form for Composition of Feeds

(Please print)

Project No. 645 Experiment No. FSA-1 Leader: Prof. A.K. Jafri

Laboratory name: Fish Nutrition Research Laboratory

Address: Department of Zoology,
Aligarh Muslim University, Aligarh, India

Laboratory director

Laboratory sample No. 8

Origin and Name of Sample

Date originally collected: March Day Monday Year 1987

Country: Northern India

State, province or department: India

County or district: U.P.

Literature reference: ICAR-Handbook of Agriculture

51

Class: Dry forage (cut and cured)

Pasture (including range plants)

Cut and fed green

Stilage

Scientific name: Genus: Calanus

Species: Calan

Variety or kind: T-21, Pant-A3

Common name of feed: Arhar

Part of plant, animal or feed product: Seed (Dust)

Process undergone before fed to animal: Drying, grinding, sieving

Stage of maturity: 150 days

Cutting or crop: 1

Height when cut (centimeters): 0150

Official grade or No.:

Fertilizer: kg per hectare

Days between last application and harvest

Purity %:

Soil classification: Alluvium

Surface texture: Loamy

Digestibility Trial

Animal: Kind: Fish (Labeo rohita)

Breed:

Sex:

Age: Years: 0 Months: 10 Weeks:

Number of animals in treatment: 10

Total number of determinations per feed per animal: 2

Average weight of animals, kg: 24.7

Physiological state: non-pregnant

pregnant 1st 2/3

fastening

losing wt

very thin ☐ thin ☐ thrifty ☐ fat ☐ very fat ☐

Percent of test ingredient in ration fed (100.0% dry matter):

Ad Libitum feeding ☒ Controlled feeding ☐

Feed fed alone ☒ Feed not fed alone, digestion by difference ☐

Method: Total feces collection ☐ feces indicator ☒

Physical form of feed: Moist cake.

Basis for calculating level of feeding: NE ME DE TDN

Length of trial: Preliminary days 03 Collection days 03

Daily dry matter consumed g kg

Daily dry matter offered g kg

Dailyorts or weighback g kg

Temperature: Degrees C (indicate if negative) 026

Pressure: millimeters

Humidity: percent

Card 40

Dry Matter

Dry matter of sample on a fed basis 101

Dry Matter: Method of analysis, if any, or none by another method but under other analyses

Anal. code factor

Digestion coeff. %

91.8

Dry Matter Basis on Which Analytical Data are Reported on This Form

NOTE: All analytical data on this sheet, except dry matter, must be expressed as indicated: a. on fed, partially dry, dry (100.0% dry matter). Where analytical data for one feed is partly on one basis and partly on another, use separate sheet for each basis or convert to same basis.

Fill in one only

Check one only

Basis: ☒ As fed ☐ Partially dry ☐ Dry (100.0% dry matter)

Proximate Principles

Ash 105 05.4

Crude fiber 106 25.7

Ether extract 107 02.4

Nitrogen-free extract 108 4.42

Protein 109 13.9

Nitrogen 212 do

Nitrogen factor 213 Write in factor to convert to protein

Energy

Gross energy 421 3800

Digestible 422

Metabolizable N-equilibrium metabolizable 423

NE_m 424

NE gain 426

NE lactating cows 427

TDN 429

Method of analysis, if any, or none by another method but under other analyses

Anal. code factor

Digestion coeff. %

05.4

25.7

02.4

4.42

13.9

do

Write in factor to convert to protein

3800

Bomb calorimeter

Harris Natl Acad Sci Natl Res Council pub 1411 1965

do

do

Lalgreen J. Animal Sci 27:793, 1968

do

Moore J. Dairy Sci 52:978

Biological

Table XV

International Source Form for Composition of Feeds

(2 parts of ml)

Project No. 645 Experiment No. FSA-1 Investigator Prof. A.K. Jafri

Card 10

12

17

22

27

32

37

42

47

52

57

62

67

72

77

82

87

92

97

102

107

112

117

122

127

132

137

142

147

152

157

162

167

172

177

182

187

192

197

202

207

212

217

222

227

232

237

242

247

252

257

262

267

272

277

282

287

292

297

302

307

312

317

322

327

332

337

342

347

352

357

362

367

372

377

382

387

392

397

402

407

412

417

422

427

432

437

442

447

452

457

462

467

472

477

482

487

492

497

502

507

512

517

522

527

532

537

542

547

552

557

562

567

572

577

582

587

592

597

602

607

612

617

622

627

632

637

642

647

652

657

662

667

672

677

682

687

692

697

702

707

712

717

722

727

732

737

742

747

752

757

762

767

772

777

782

787

792

797

802

807

812

817

822

827

832

837

842

847

852

857

862

867

872

877

882

887

892

897

902

907

912

917

922

927

932

937

942

947

952

957

962

967

972

977

982

987

992

997

1002

1007

1012

1017

1022

1027

1032

1037

1042

1047

1052

1057

1062

1067

1072

1077

1082

1087

1092

1097

1102

1107

1112

1117

1122

1127

1132

1137

1142

1147

1152

1157

1162

1167

1172

1177

1182

1187

1192

1197

1202

1207

1212

1217

1222

1227

1232

1237

1242

1247

1252

1257

1262

1267

1272

1277

1282

1287

1292

1297

1302

1307

1312

1317

1322

1327

1332

1337

1342

1347

1352

1357

1362

1367

1372

1377

1382

1387

1392

1397

1402

1407

1412

1417

1422

1427

1432

1437

1442

1447

1452

1457

1462

1467

1472

1477

1482

1487

1492

1497

1502

1507

1512

1517

1522

1527

1532

1537

1542

1547

1552

1557

1562

1567

1572

1577

1582

1587

1592

1597

1602

1607

1612

1617

1622

1627

1632

1637

1642

1647

1652

1657

1662

1667

1672

1677

1682

1687

1692

1697

1702

1707

1712

1717

1722

1727

1732

1737

1742

1747

1752

1757

1762

1767

1772

1777

1782

1787

1792

1797

1802

1807

1812

1817

1822

1827

1832

Table XVI

International Source Form

(Please print)

Project No. 645 Experiment No. FSA-1 Leader Prof. A.K. Jafri

Laboratory name Fish Nutrition Research Laboratory

Address Department of Zoology, Aligarh Muslim University, Aligarh, India

Laboratory director

Laboratory sample No. 10

Origin and Name of Sample March Monday 1987

Date originally collected Month Day Year

Country Northern India

State, province or department India

County or district Aligarh

Literature reference ICAR-Handbook of Agriculture

Class Dry forage (cut and cured)

Pasture (including range plants)

Cut and fed green

Silage

Scientific name: Genus Triticum

Species aestivum

Variety or kind Sonalika

Common name of feed Gehun, Gam

Part of plant, animal or feed product Grain (Bran)

Process undergone before fed to animal Drying, grinding, sieving

Stage of maturity 90 days

Cutting or crop 1

Height when cut (centimeters) 0 0 8 0

Official grade or No. 16

Fertilizer: kg per hectare N 0 0 7 0 P 0 0 3 5 K 0 0 2 2

Days between last application and harvest

Purity %

Chief contaminant

Soil classification Alluvium

Surface texture Loamy

Digestibility Trial

Animal: Kind Fish (Labeo rohita)

Breed

Sex

Age: Years 0 + Months Weeks

Number of animals in treatment 10

Total number of determinations per feed per animal 2

Average weight of animals, kg or 9 2 4 7

Physiological state: non-pregnant pregnant 1st 2/3 pregnant last 1/3

losing wt maintaining wt gaining wt fattening

very thin thin thrifty fat very fat

Percent of test ingredient in ration fed (100.0% dry matter)

Ad Libitum feeding

Feed fed alone

Method: Total feces collection feces indicator

Physical form of feed Moist cake.

Basis for calculating level of feeding: Preliminary days 0 3 Collection days 0 3

Length of trial: kg kg kg

Daily dry matter consumed 9

Daily dry matter offered 9

Dailyorts or weighback 9

Temperature: Degrees C (indicate if negative) 0 2 6

Pressure: millimeters percent

Humidity: Card 40

Dry Matter

Dry matter of sample on an as fed basis

Method of analysis: if analysis done by another method put under other analysis

Anal. code factor

Digestion coeff %

Dry Matter Basis on Which Analytical Data are Reported on This Form

NOTE: All analytical data on this sheet, except dry matter, must be expressed on indicated basis: as fed, initially dry, dry (100.0% dry matter). Where analytical data for one feed is partly on one basis and partly on another use separate sheet for each basis or convert to same basis

Check one only

Basis

As fed

Partially dry

Dry (100.0% dry matter)

Proximate Principles

Ash

Crude fiber

Ether extract

Nitrogen-free extract

Protein

Nitrogen

Nitrogen factor

Energy

Gross energy

Digestible

Metabolizable

N-equilibrium metabolizable

NE_m

NE gain

NE lactating cows

TDN

Biological

%

Read instructions before filling in form.

Food and Agriculture Organization of the United Nations (FAO, AGA)

00100 Rome, Italy

Table XVII

International Source Form for Composition of Feeds

(Please print)

Project No.	645	Experiment No.	FSA-1	Leader	Prof. A.K. Jafri	Card 10	12	17	22
Laboratory name	Fish Nutrition Research Laboratory								
Address	Department of Zoology, Aligarh Muslim University, Aligarh, India								
Laboratory director									
Laboratory sample No.	11								
Origin and Name of Sample	Date originally collected Month March Day Monday Year 1987								
Country	India								
Region	Northern India								
State, province or department	U.P.								
County or district	Aligarh								
Literature reference	ICAR-Handbook of Agrilculture 51								
Class	Dry forage (cut and cured)	Pasture (including range plants)	Cut and fed green	Silage					
Scientific name: Genus	ORYZA								
Species	SATIVA								
Variety or kind	IR-8								
Common name of feed	Dhan, Chawal								
Part of plant, animal or feed product	Grains (Polish, Bran)								
Process undergone before fed to animal	Drying, grinding, sieving								
Stage of maturity	85-90 days								
Cutting or crop	1	Height when cut (centimeters)	0065	12					
Official grade or No.	16	20	24	28					
Fertilizer: kg per hectare	N	0090	P	0045	K				
Days between last application and harvest									
Purity %									
Chief contaminant									
Soil classification	Alluvium								
Surface texture	Clayey loam								
Digestibility Trial									
Animal: Kind	Fish (Labeo rohita)								
Breed	Sex								
Age: Years	0	Months	Weeks						
Number of animals in treatment	10								
Total number of determinations per feed per animal	2								
Average weight of animals, kg	or g 24.7								
Physiological state: non-pregnant	pregnant 1st 2/3								
losing wt	maintaining wt								
	gaining wt								
	fattening								

Food Information Center
of the United Nations (FAO, AGA)
Via delle Terme di Caracalla
00100 Rome, Italy

Read instructions before filling in form.

very thin	thin	thrifty	fat	very fat
Percent of test ingredient in ration fed (100.0% dry matter)				
Ad Libitum feeding	<input checked="" type="checkbox"/>	Controlled feeding		
Feed fed alone	<input checked="" type="checkbox"/>	Feed not fed alone, digestion by difference		
Method: Total feces collection		Feces indicator	<input checked="" type="checkbox"/>	

Physical form of feed Moist cake.

Basis for calculating level of feeding:	NE	ME	DE	TDN
Length of trial: Preliminary days	03	Collection days	03	
Daily dry matter consumed	g	kg		
Daily dry matter offered	g	kg		
Dailyorts or weightback	g	kg		
Temperature: Degrees C (indicate if negative)	0	2	6	
Pressure: millimeters				
Humidity: percent				

Card 40

Dry Matter

Dry Matter %	Method of analysis, if analysis is done by another method put under other analyses	Anal. code factor	Digestion coeff %
96.7	Above 105°C or in vacuum		

Dry Matter Basis on Which Analytical Data are Reported on This Form

NOTE: All analytical data on this sheet, except dry matter, must be expressed as indicated (i.e., as fed, partially dry, dry (100.0% dry matter). Where analytical data for one feed is partly on one basis and partly on another use separate sheet for each basis or convert to same basis.

Fill in one only

Check one only	102	103	104
As fed	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Partially dry	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Dry (100.0% dry matter)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Proximate Principles

Ash	105	106	107	108	109	212	213
Crude fiber	14.4	Wende	Assoc. Official Agr. Chem. 1960				
Ether extract	01.8	Assoc. Official Agr. Chem. 1960					
Nitrogen-free extract	05.8	By difference					
Protein	62.5	Kjeldahl					
Nitrogen		do					
Nitrogen factor		Write in factor to convert to protein					

Energy

Gross energy	421	Bomb calorimeter		
Digestible	422	Harris Natl. Acad. Sci. Natl. Res. Council pub. 1411 1966		
Metabolizable	423	do		
Non-equilibrium metabolizable	424	do		
NE _m	426	Loggreen J. Animal Sci. 27:793. 1968		
NE gain	427	do		
NE lactating cows	430	Moore J. Dairy Sci. 52:978		

%

TDN		Biological	
-----	--	------------	--

Table XVIII

International Source Form for Composition of Feeds

Project No. 645 Experiment No. FSA-1 Investigator Prof. A.K. Jafri

Laboratory name Fish Nutrition Research Laboratory

Address Department of Zoology,
Aligarh Muslim University, Aligarh, India

Laboratory director

Laboratory sample No.

12

Origin and Name of Sample

Date originally collected Month March Day Monday Year 1987

Country

India

Region Northern India

Altitude (meters)

36

State, province or department

U.P.

County or district

Aligarh

Literature reference ICAR-Handbook of Agriculture

Class Dry forage (cut and cured) Pasture (including range plants) Cut and fed green Silage

Scientific name : Genus Hordeum

Species vulgare

Variety or kind Kailash

Common name of feed Jow

Part of plant, animal or feed product Grains (Dust)

Process undergone before fed to animal Drying, grinding, sieving

Stage of maturity 100 days

Cutting or crop 1 Height when cut (centimeters) 0 19 10

Official grade or No.

Fertilizer: kg per hectare

Days between last application and harvest

Purity % Chief contaminant

Soil classification Alluvium

Surface texture Moderately rich loam

Digestibility Trial

Animal: Kind Fish (Labeo rohita)

Breed Sex

Age: Years 0 Months 10 Weeks

Number of animals in treatment 10

Total number of determinations per feed per animal 2 period No.

Average weight of animals, kg or g 2 4 7

Physiological state: non-pregnant pregnant 1st 2/3 pregnant last 1/3

losing wt maintaining wt gaining wt fattening

very thin	thin	thrifty	fat	very fat
Percent of test ingredient in ration fed (100.0% dry matter)				
Ad Libitum feeding	<input checked="" type="checkbox"/>	Controlled feeding		
Feed fed alone	<input checked="" type="checkbox"/>	Feed not fed alone, digestion by difference		
Method: Total feces collection		feces indicator	<input checked="" type="checkbox"/>	

Physical form of feed Moist cake.

Basis for calculating level of feeding:	NE	ME	DE	IDN
Length of trial: Preliminary days	0 3	Collection days	0 3	
Daily dry matter consumed	g	kg		
Daily dry matter offered	g	kg		
Dailyorts or weightback	g	kg		
Temperature: Degrees C (indicate if negative)	0 2 6			
Pressure: millimeters				
Humidity: percent				
Card 40				

Dry Matter

Dry matter of sample on fed basis

8 9 8

Dry Matter: Method of analysis, if analyzed by another method put under analysis factor

Dry Matter Basis on Which Analytical Data are Reported on This Form

NOTE: All analytical data on this sheet, except dry matter must be expressed as indicated: a. as fed, normally dry (100.0% dry matter) Where analytical data for one feed is partly on one basis and partly on another use separate sheet for each basis or convert to same basis

Proximate Principles

Ash	105	0 3 5	Assoc Official Agr Chem 1960		
Crude fiber	106	0 8 6	Weende		
Ether extract	107	0 0 6	Assoc Official Agr Chem 1960		
Nitrogen-free extract	108	6 9 2	By difference		
Protein	109	0 7 9	Kjeldahl		
Nitrogen	212		do		
Nitrogen factor	213	6 2 5	Write in factor to convert to protein		

Energy

Gross energy	421	4 0 3 6	Bomb calorimeter		
Digestible	422		Harris Natl Acad Sci Natl Res Council pub 1411 1966		
Metabolizable	423		do		
N-equilibrium metabolizable	424		do		
NE _m	426		Lofgren J Animal Sci: 27:793 1968		
NE gain	427		do		
NE lactating cows	430		Moe J Dairy Sci: 52:978		

TDN

Biological

CHAPTER - II

ENERGY REQUIREMENT OF THE INDIAN MAJOR CARP, LABEO ROHITA (HAM)

INTRODUCTION

Livestock and poultry nutritionists have long recognised the importance of meeting the energy requirements of animals in formulating practical diets. Various feeding tables listing the allowance for energy and other nutrients are available for the farm animals. Relatively little information is available on the energy requirements of fish (Adron et al., 1976; Marries and Klissil, 1979; Cho and Kaushik, 1985; Machiels and Henken, 1985; Daniels and Robinson, 1986; Barrows et al., 1988; and Colosso et al., 1988). Fish nutritionists have generally given priority to meeting the requirements for protein, minerals and vitamins, although it has been recognised that deficient energy level in the ration may lead to less efficient utilization of protein and other nutrients by the fish. Intake of higher than the required energy may cause excessive fat deposition (Adron et al., 1976; Garling and Wilson, 1976, and Takeuchi et al., 1978a,b,c).

Fish, like other animals, are known to eat to meet their energy requirements (Lee and Putnam, 1973; and Page and Andrews, 1973) and thus have the capability of compensating for a low energy density in food by eating more of it, provided the food is nutritionally balanced. Such a compensation, however, occurs up to the physical limit of the digestive tract, although a high energy density diet will require less feed per unit of weight gained. However, maximal physical intake of high energy density diet sustains a higher growth rate. Provision of an optimal balance of energy components in diet is, therefore, important from the view point of fish production and the cost-effectiveness of the culture system.

Fish are ammoniotelic animals and can readily use protein as an energy source. In the event of disproportionately high protein to non-protein energy ratio in diet, greater utilization of protein may occur, leading to an excessive concentration of ammonia in the culture system (Beamish and Thomas, 1984).

A deficiency of non-protein energy sources (lipid and carbohydrate) in diet may retard growth rate (Lovell, 1976), since protein will then be used more for energy purposes rather than for protein synthesis. Similarly, if the diet contains an excess of non-protein energy, appetite or demand may be

satisfied before a sufficient quantity of protein (and also other nutrients) is ingested to satisfy demand for protein synthesis and growth. In an exhaustive review on protein requirements of fish, Wilson and Halver (1986) emphasized that dietary protein requirement for fish may also be influenced by the dietary protein to energy balance. Estimation of protein energy ratio in the diet for optimum growth of the fish is thus an important aspect in fish nutritional studies.

Besides protein to energy ratio in diet, several other factors may influence the energy requirement of fish. It has been pointed out that, in identical condition, the energy requirement varies with the species (Beamish, 1964; Dean and Goodnight, 1964; and Brett and Groves, 1979). Temperature, by altering the kinetics of metabolic reaction, has a direct influence on the energy requirement, as measured by increased or decreased growth rates, respiratory or oxygen consumption (Rozin and Mayer, 1961; Dean and Goodnight, 1964; and Brett, 1965). The energy requirement is also reported to be size and/or age dependent (Brown, 1957; Brett et al., 1969; Pandian, 1967, 1970; de Silva and Balbontin, 1974).

Several studies have been made in the past to assess the effect of total energy intake on the growth

and conversion efficiencies of different species of fish. Although the gross protein requirement of the Indian major carp, L. rohita, has been reported in the past (Sen et al., 1978; and Renukaradhya and Varghese, 1986), no information is available on the energy requirements of this species. Intensification of carp aquaculture would require the use of well-balanced formulated feed as per the requirement of the species. The lack of data on complete nutritional requirements of Indian major carp is one of the major technical constraints in the development of least-cost and nutritionally adequate production diet for this group of fish.

The present study deals with the effect of different energy levels on the growth and conversion efficiencies, leading to optimization of an energy: protein ratio in L. rohita. Knowledge of energy requirement of this fish at a given protein level would be useful in identifying the best calorie-to-protein ratio, as also in balancing the energy and protein components, while formulating practical diets for the above species.

MATERIALS AND METHODS

I. Source of fish and acclimatization of test lots

Fingerlings of L. rohita were obtained from the Uttar Pradesh Fish Seed Centre, Kolahar, Mathura. These were transported to the laboratory in oxygen filled polythene bags and stocked in earthen fish pond (12 x 6 x 1.5 m). Supplementary feeding at the level of 5% of the biomass was carried out using the conventional, rice bran: mustard oil cake (1 : 1) feed mixture, daily at 0800 h. The required number of fingerlings were transferred to the wet laboratory, disinfected with 5 ppm KMnO_4 solution and acclimatized on casein-gelatin test diet for about two weeks.

II. Preparation of experimental diets

The reported gross protein requirements of L. rohita is around 40-45% of the diet (Sen et al., 1978; Renukaradhya and Varghese, 1986). Based on these finding, three approximately iso-nitrogenous diets were prepared, using casein and gelatin as the protein source (Mertz, 1972), to work out the energy requirement of this species. The energy level in the experimental diets ranged from 3390 - 3870 cal/g dry

diet (Table I). The mineral and vitamin premixes (Table II,III) were prepared according to Halver (1976). The protein to energy (P/E) ratios in the experimental diets ranged from 75.1 - 86.0 kcal/mg protein.

The calculated quantities of various dietary ingredients were weighed on an electronic balance. A known quantity of water was then taken in a stainless steel attachment of Hobart electric mixer and heated to 80°C. Gelatin was dissolved into it with slow stirring and heating the content on to a double boiler unit. The mixer bowl was removed from heating, dextrin and mineral mix added to it, and the content blended in Hobart mixer while still in lukewarm state. This was followed by the addition of remaining ingredients. As the diet began to solidify, the speed of stirring was increased gradually to incorporate air into the mixture. The final diet, about the consistency of bread dough, was poured into a teflon-coated pan and placed in a refrigerator to jell.

For the preparation of crumbles, the above material was cold extruded through Hobart extruder fitted with 1 mm die. The strands were spread over the receiving tray and placed in hot air oven at 60°C. The diet upon drying were crumbled and sieved through ASTM-18 and ASTM-30 screens, and stored in sealed plastic bags in a freezer until used.

III. Feeding trial

Fingerlings (2.9-5.3 cm total length) were sorted out from the acclimatized fish lot maintained in the wet laboratory and stocked in 70 l high density polyvinyl circular troughs (water volume 55 l) fitted with a continuous water flow-through system. The stocking density was 100 fish per trough. The water exchange rate in each trough was maintained at 1-1.5 l/min. The bottom of the trough was siphoned off every morning to remove unused food, if any, and the faecal matter. A natural light : dark cycle was maintained.

The fish were fed test diet crumbles at the rate of 3% body weight daily at 0800 and 1600 h, except on the day when weekly measurements were taken. The feeding level and feeding schedule were chosen after carefully observing the feeding behaviour of the fish and their dietary intake. Feeding allowances were adjusted on weekly basis. Care was taken to ensure that diet crumbles were quickly consumed. Accumulation of diet at the bottom was avoided to prevent any possible leaching of the nutrients. A thorough scrubbing and cleaning of each trough was also carried out weekly. The average water temperature and dissolved oxygen over the 6-week feeding

trial, based on daily measurements, ranged between 24-28°C and 6.3 - 6.8 ppm, respectively.

Survival was 100% in all the dietary treatments.

IV. Assessment of nutritional parameters

One of the criteria chosen to ascertain the performance of feed was the assessment of growth obtained by subtracting the initial weight of biomass from its final weight over a particular duration of feed trial.

The specific growth rate (%) was evaluated using the formula:

$$\text{SGR (\%)} = \frac{W_2 - W_1}{D} \times 100$$

Where W_1 = Initial weight of biomass (g)

W_2 = Final weight of biomass (g)

D = Duration of feed trial (days)

The food conversion ratio (FCR) was calculated on the basis of the ratio of food consumed to live weight gained by the fish. The gross growth efficiency represented the ratio of live weight gain to food consumed. The protein efficiency ratio (PER) was calculated on the basis of the ratio of protein consumed to live weight gained.

RESULTS

Results of the 6-week feeding trial conducted on L. rohita to quantify its energy requirement has been summarized in Table IV.

The average weight of fish over the experimental period for the test diets used increased linearly on their initial weight (Fig. I). The fish registered the maximum gain in live weight with 3500 cal/g dry diet energy level. The lowest gain was recorded with 3870 cal/g dietary energy. The weekly mean live weight gain per cent for different treatments have been plotted in Fig. II . In terms of percentage, the maximum (89%) gain, occurred in fish fed 3500 cal/g energy diet, whereas it was only 72% at high (3870 cal/g) dietary energy intake.

The average specific growth rate (%) of fish at various energy intakes have been depicted in Fig. III. The specific growth rate increased linearly with increasing dietary energy content up to the incorporation rate of 3500 cal/g and declined sharply at 3870 cal/g energy intake.

The FCR and gross growth efficiency in fish fed different levels of energy have been shown in Fig. IV . Marked differences were seen in the food

conversion values. The best (2.1) FCR was noted at 3500 cal/g dietary energy level. Less efficient conversion was obtained with diets having levels of higher or lower to the requirement (3500 cal/g). The gross growth efficiency and the PER were likewise maximum, 0.48 and 1.15, respectively, at the above level of dietary energy.

Fat and water contents of the carcass was found affected with the levels of energy intake (Table IV). These two constituents also indicated a negative relationship with each other. Highest (32% dry weight) amount of crude fat, noted in the carcass of fish fed diet containing 3870 cal/g energy, was accompanied with lowest percentage of water.

DISCUSSION

The maintenance energy needs of fish are considerably lower than those of homeothermic animals (Cho and Kaushik, 1985). However, compared to higher vertebrates, usually a high proportion of heat production is derived from dietary protein. Although the utilization of protein for basal energy metabolism is an established phenomenon in fish, conventional energy yielding nutrients like fats and carbohydrates can reduce the oxidation of protein to satisfy the

energy needs of fish and thus improve the utilization of dietary protein for growth (Cowey, 1975; and Watanabe, 1977). The beneficial effects of incorporating such protein-sparing nutrients have been widely studied (Lee and Putnam, 1973; Takeuchi et al., 1978 a,b,c; 1979; Bromley, 1980; Millikin, 1983; and Beamish and Medland, 1986), and optimal ratios between protein and energy proposed for many species of fish (Cho and Kaushik, 1985).

It would appear, both from the plot of average fish weight against time as well as the pattern of changes observed in the specific growth rate, that the level of energy producing the maximum growth in L. rohita is around 3500 cal/g. However, further increase of dietary energy intake to 3870 cal/g resulted in smaller gain than that observed at the low energy (3390 cal/g) intake. The findings are in conformity to the results reported by Barrows et al. (1988) on Walleye fingerlings. Takeuchi et al. (1978 b) and Millikin (1983) noticed similar effects of dietary energy on weight gains in rainbow trout and striped bass, respectively.

The results of the present study indicates that a proper balance between energy and protein is important for optimum growth and feed efficiency of L. rohita, as is true for other fish species. The

energy to protein ratio of 83.3 (kcal/mg protein) recorded for the maximum growth of this species is identical to the value reported for small-sized common carp (Takeuchi et al., 1978d and Lovell, 1984b). Garling and Wilson (1977), using purified diets, reported the optimum energy to protein ratio for the growth of young channel catfish to be 96, while Page and Andrews (1973), using practical diets, found the value to be 97. Stickney and Lovell (1977), on the other hand, recommended energy requirement of 80 to 90 kcal/mg of protein for the fingerling and production diets of the above species. Higher values reported for the channel catfish may be attributed to the fact that the fish, being carnivore, requires higher amount of energy per unit weight of protein.

In growing animals, part of the retained energy is stored as protein and part as fat. Besides water temperature and maturity stages, deposition of these energy resources depends upon the amount by which the dietary energy intake exceeds the energy expended as heat. In almost all cases retention of energy and deposition of new tissue result in an increase in weight of the animal. High energy diets, especially when a higher percentage of the calories is from fat, will produce fatty fish which have a lower dressing percentage than fish fed a low energy diets. In the present

study the diet containing minimum carbohydrate and maximum amount of fat as well as highest energy (3870 cal/g) was found to produce fatty (32% dry weight) carcass. Enhanced lipid deposition noted in L. rohita with increased energy intake was in agreement with observations on other fish species (Cowey and Sargent, 1979; Watanabe, 1982; Jobling and Wandsvik, 1983; Machiels and Henken, 1985; Daniels and Robinson, 1986; and Barrows et al., 1988). However, additional feed energy was not reported to increase the fat content of the dry carcass in brook trout (Ringrose, 1971). The observed negative relationship between fat and water content of the body of L. rohita was similar to the findings on several other fish species (Cowey, et al., 1972; Austreng and Refstie, 1979; Reinitz and Hitzel, 1980; and Watanabe, 1982).

Decreased PER and FCR noted during the present study with higher dietary energy was also seen in juvenile red drum (Daniels and Robinson, 1986).

SUMMARY

The effect of different energy levels on the growth and conversion efficiencies, leading to optimization of energy: protein ratio, has been reported in the fingerlings of the Indian major carp, L. rohita.

The best growth, food conversion and protein efficiency were noted with diet containing 3500 cal/g of energy, corresponding to 83.3 kcal/mg protein. Intake of dietary calorie below and above the aforesaid level resulted in poor growth. Moisture and fat constituents of carcass were also found to be influenced by varying calorie intake. Maximum fat deposition was observed in fish fed diet containing the highest (3870 cal/g) energy. The study indicates that, for practical diet formulation for this species, incorporation of 3500 cal/g of energy is desirable for optimum fish production.

Table I Ingredient composition of the test diets
used for the estimation of energy require-
ment

Ingredients g/100g diet	Dietary protein level		
	45%	42%	45%
Casein	40.17	38.00	40.17
Gelatin	12.84	12.00	12.84
Dextrin	28.00	28.00	18.99
✓-cellulose	7.99	8.00	8.00
Corn oil	4.00	6.00	10.00
Cod liver oil	2.00	3.00	5.00
Mineral mix	4.00	4.00	4.00
Vitamin mix	1.00	1.00	1.00
Water	100.00	100.00	100.00
Gross energy (kcal/100g dry diet) *	338.00	350.00	387.00
Protein : energy ratio (kcal/mg protein)	75.00	83.30	86.00

*Calculated on the basis of physiological fuel value,
4.5, 8.5 and 3.0 kcal/g of protein, fat and carbohydrate,
respectively.

Table II Mineral premix

	g/100g
Calcium biphosphate	13.48
Calcium lactate	32.40
Ferric citrate	2.97
Magnesium sulphate	13.70
Potassium phosphate (dibasic)	23.86
Sodium biphosphate	8.72
Sodium chloride	4.35
Aluminium chloride 6 H ₂ O	0.015
Potassium iodide	0.015
Cuprous chloride	0.010
Manganous sulphate H ₂ O	0.080
Cobalt chloride 6 H ₂ O	0.100
Zinc sulphate 7 H ₂ O	0.300

Table III Vitamin premix

	g/100 g
α -cellulose	8.0
Choline chloride	0.500
Inositol	0.200
Ascorbic acid	0.100
Niacin	0.075
Calcium pantothenate	0.050
Riboflavin	0.020
Manadione	0.004
Pyridoxine Hcl	0.005
Thiamin Hcl	0.005
Folic acid	0.0015
Biotin	0.0005
α -tocopherol acetate	0.040
Vitamin B ₁₂ (10 mg/500 ml H ₂ O)	0.5 ml

Table IV Results of feeding purified test diets
to Labeo rohita during a 6-week feeding
trial

	DIETARY ENERGY LEVEL (kcal/100g, dry diet)		
	338.00	350.00	387.00
Initial individual weight (g)	1.10	0.98	1.01
Final individual weight (g)	1.93	1.85	1.74
Percentage increase in weight	74.93	89.08	71.72
Specific growth rate(%)	1.33	1.52	1.28
Food conversion ratio*	2.63	2.06	2.56
Protein efficiency ratio	0.85	1.15	0.86
Gross growth efficiency	0.38	0.48	0.39
Percentage survival	100	100	100
Carcass Composition			
Moisture (%)	70.75	70.29	68.18
Fat (% dry wt. basis)	25.43	28.29	32.15

* Dry to wet basis

Results are mean of duplicate runs

Fig. I : Weekly changes in average fish weight (g) at
varying energy levels (●—● 338 kcal; ○—○
350 kcal; ▲—▲ 387 kcal).




Fig. II : Growth of L. rohita on various dietary energy levels (●—● 350 kcal; ○—○ 387 kcal; ▲—▲ 338 kcal).




Fig. III : Specific growth rate (%) of L. rohita fed
varying energy levels.





Fig. IV : Food conversion ratio (●—●) and gross growth efficiency (O-----O) of L. rohita fed varying energy levels.



REFERENCES

- Abalos, T.U., 1989 The effect of live maggots and growth hormone in practical feeds on growth of tilapia (Oreochromis niloticus) fingerlings. In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst.).
- Adron, J.W., 1976 Effects of dietary energy level and dietary energy source on growth, feed conversion and body composition of turbot (Scophthalmus maximus L.). Aquaculture 7: 125-132.
- Blair, A.,
Cowey, C.B. and
Shanks, A.M.
- A.O.A.C. 1984 Official Methods of Analysis, 14th ed. Association of Official Analytical Chemists, Washington, D.C.
- Austreng, E. and 1979 Effects of varying dietary protein level in different families of rainbow trout, Aquaculture 18: 145-156.
- Refstie, T.
- Austreng, E., 1980 Digestibility of fat and fatty acids in rainbow trout and minck. Aquaculture 19: 93-95.
- Skrede, A. and
Eldegard, A.
- Barrows, F.T., 1988 Effects of dietary protein and energy levels on weight gains, body composition, and RNA:DNA ratios of fingerling Walleyes. Prog. Fish Cult. 50: 211-218.
- Sell, J.L. and
Nickum, J.G.
- Beamish, F.W.H. 1964 Respiration of fishes with special emphasis on standard oxygen consumption. Influence of weight and temperature on respiration of several species Can. J. Zool. 42: 177-188.
- Beamish, F.W.H. 1986 Protein sparing effects in large rainbow trout, Salmo gairdneri. Aquaculture 55: 35-42.
- and Medland, T.R.

- Beamish, F.W.H. and Thomas, E. 1984 Effects of dietary protein and lipid on nitrogen losses in rainbow trout, Salmo gairdneri. Aquaculture 41: 359-371.
- Bergot, F. 1979a Carbohydrate in rainbow trout diets: Effects of the level and sources of carbohydrate and number of meals on growth and body composition. Aquaculture 18: 157-167.
- Bergot, F. 1979b Problèmes particuliers posés par l'utilisation des glucides chez la truite arc-en-ciel. Ann. Nutr. Aliment. 33: 247-257.
- Bhuiyan, A.K.M.A., Begum, N.N., Begum, M. and Haq, M.E. 1989 Survey of potential fish feed ingredients of Bangladesh. In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst).
- Brett, J.R. 1965 The relation of size to rate of oxygen consumption and sustained swimming speed of sockeye salmon (Oncorhynchus nerka). J. Fish. Res. Bd. Can. 22: 1491-1501.
- Brett, J.R. and Groves, T.D.D. 1979 Physiological energetics. In: Fish Physiology. Vol. VIII. Academic Press, London/New York (eds. Hoar, W.S., Randall, D.J. and Brett, J.R.). pp. 280-352.
- Brett, J.R., Shelbourn, J.E. and Shoop, C.T. 1969 Growth rate and body composition of fingerling sockeye salmon, Oncorhynchus nerka, in relation to temperature and ration size. J. Fish. Res. Bd. Can. 26: 2363-2394.
- Bromley, P.J. 1980 Effect of dietary protein lipid and energy content on the growth of turbot (Scophthalmus maximus L.). Aquaculture 19: 359-369.

- Brown, M.E. 1957 Experimental studies on growth. In: The Physiology of Fishes. Vol. 1. Academic Press, New York (ed. Brown, M.E.). pp. 361-400.
- Cagauan, A.G. and Pullin, R.S.V. 1989 Azolla in Aquaculture: Past, present and future. In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst).
- Castell, J.D. 1979 Review of lipid requirements of finfish. In: Finfish Nutrition and Fishfeed Technology Vol. I, Heenemann, Berlin, Federal Republic of Germany (eds. Halver, J.E. and Tiews, K.). pp. 59-84.
- Chakrabarty, R.D., Sen, P.R., Chatterjee, D.K. and Kowtal, G.V. 1973 Observations on the relative usefulness of different feed for carp spawn and fry. J. Inland, Fish. Soc. Ind. 5: 182-188.
- Chattopadhyay, N.R. and Konar, S.K. 1987 Use of groundnut oil cake in rearing of Indian major carp spawn. In: The First Asian Fisheries Forum, Mangalore, Karnataka, Dec. 4-8, 1987 (Abst).
- Chiayvareesajja, S., Wongwit, C. and Tansakul, R. 1989 Cage culture of tilapia (Oreochromis niloticus) using aquatic weed mixture pellet. In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst).
- Cho, C.Y., Bayley, H.S. and Slinger, S.J. 1974 Partial replacement of herring meal with soybean meal and other changes in diet for rainbow trout (Salmo gairdneri). J. Fish. Res. Bd. Can. 31(9): 1523-1528.
- Cho, C.Y., Cowey, C.B. and Watanabe, T. 1985 Finfish Nutrition in Asia: Methodological approaches to research and development. International Development Research Centre, Ottawa, Canada.

- Cho, C.Y. and Kaushik, S.J. 1985 Effects of protein intake on metabolizable and net energy values of fish diets. In: Nutrition and Feeding in Fish. Academic Press, London/New York. (eds. Cowey, C.B., Mackie, A.M. and Bell, J.G.). pp. 95-117.
- Chow, K.W. 1982 Carp Nutrition Research at the Freshwater Aquaculture Research and Training Centre, Dhauli (India) FI/DP/IND/75/031/Field-Doc. 4, FAO, Rome, pp. 1-30.
- Chow, C.Y. and Woo, N.Y.S. 1989 Bioenergetic studies on the growth of an omnivorous fish, Oreochromis mossambicus : Evaluation of utilization of spirulina algae in feed. In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst).
- Coloso, R.M., Benitez, L.V. and Tiro, L.B. 1988 The effect of dietary protein-energy levels on growth and metabolism of milkfish (Chanos chanos Forsskal) Comp. Biochem. Physiol. 89 A(1): 11-17.
- Cowey, C.B. 1975 Aspects of protein utilization by fish. Proc. Nutr. Soc. 34: 57-63.
- Cowey, C.B., Luquet, P. 1983 Physiological basis of protein requirements of fishes. Critical analysis of allowances. In: Protein Metabolism and Nutrition. Vol. I, INRA, Paris (eds. Pion, R., Arnal, M. and Bonin, D). pp. 364-384.
- Cowey, C.B., Pope, J.A., Adron, J.W. and Blair, A. 1972 Studies on the nutrition of marine flatfish. The protein requirement of plaice (Pleuronectes platessa). Br. J. Nutr. 28: 447-456.
- Cowey, C.B. and Sargent, J.R. 1972 Fish nutrition. Adv. Mar. Biol. 10: 383-492.

- | | | |
|---------------------------------------|------|--|
| Cowey, C.B. and Sargent, J.R. | 1979 | Nutrition. In: Fish Physiology. Vol. VIII, Academic Press, London/New York (eds. Hoar, W.S., Randall, D.J. and Brett, J.R.). pp. 1-69. |
| Daniels, W.H. and Robinson, E.H. | 1986 | Protein and energy requirements of juvenile red drum (<u>Sciaenops ocellatus</u>). <u>Aquaculture</u> 53: 243-252. |
| Dean, J.M. and Goodnight, C.J. | 1964 | A comparative study on carbohydrate metabolism in fish as affected by temperature and exercise. <u>Physiol. Zool.</u> 37: 280-299. |
| Dehadrai, P.V. and Thakur, N.K. | 1980 | Magur and Singhi Culture. Central Inland Fisheries Research Institute, Barrackpore, India, 7 p. |
| de Silva, S.S. and Balbontin, F. | 1979 | Laboratory studies on food intake, growth and food conversion of young herring, <u>Clupea harengus</u> (L). <u>J. Fish Biol.</u> 6: 645-658. |
| Dupree, H.K. and Sneed, K.E. | 1966 | Response of channel catfish fingerlings to different levels of major nutrients in purified diets. US. Bur. Sport. Fisheries Wildl. Tech. Pap. 9: 1-21. |
| Ensminger, M.E. and Olentine, Jr.C.G. | 1980 | Feed and Nutrition. Ensminger Publishing Company, Clovis, California. U.S.A. 1417 p. |
| FAO/UNDP | 1979 | Report of the consultancy on the fish nutrition programme at the Changi Marine Fisheries Research Centre, Primary Production Department, Singapore. SCS/79/WP/85, New M.B. FAO/UNDP. South China Fish. Dev. Coord. Prog., Manila, Philippines, 25 p. |


- | | | |
|--|------|---|
| Garling, D.L. Jr. and
Wilson R.P. | 1976 | The optimum dietary protein to energy ratio for channel catfish fingerlings, <u>Ictalurus punctatus</u> . <u>J. Nutr.</u> <u>106</u> : 1368-1375. |
| Garling, D.L. Jr. and
Wilson, R.P. | 1977 | Effect of dietary carbohydrate to lipid ratios on growth and body composition of fingerling channel catfish. <u>Prog. Fish Cult.</u> <u>39</u> : 43-47. |
| Gatlin, D.M. and
Stickney, R.R. | 1982 | Fall winter growth of young channel catfish in response to quantity and source of dietary lipid. <u>Trans. Am. Fish. Soc.</u> <u>111</u> : 90-93. |
| Hajra, A. | 1987 | Biochemical investigations on the protein-calorie availability in grass carp (<u>Ctenopharyngodon idella</u> Val.) from an aquatic weed. (<u>Ceratophyllum demersum</u> Linn.) in the tropics. <u>Aquaculture</u> <u>61</u> (2): 113-120. |
| Halinge, N.V.,
Shetty, H.P.C. and
Nandeesh, M.C. | 1989 | Evaluation of three terrestrial fodder grasses as feed for the grass carp, <u>Ctenopharyngodon idella</u> (Valenciennes). In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst). |
| Halver, J.E. | 1976 | Nutritional deficiency diseases in salmonids. <u>Fish. Pathology</u> <u>10</u> (2): 165-180. |
| Halver, J.E. | 1979 | The nutritional requirements of cultivated warm-water and coldwater fish species. In: Advances in Aquaculture, Fishing News Books Ltd., Farnham, Surrey, England (eds. Pillay, T.V.R. and Dill, Wm. A.). pp. 574-580. |

- Halver, J.E. 1982 Fish nutrition and diet development. A report prepared for the development of intensive freshwater fish culture project. FAO/FI/DP/Hun/79/001/Field-dec-3. FAO. Rome.
- Halver, J.E. 1985 Recent advances in vitamin nutrition and metabolism in fish. In: Nutrition and Feeding in Fish. Academic Press, London/New York (eds. Cowey, C.B., Mackie, A.M. and Bell, J.G.). pp. 415-429.
- Haque, N.S. 1988 Protein, energy and nucleic acid levels in catfish, Heteropneustes fossilis fed with four diets. In: National Symposium of Aquaculture Productivity, New Delhi, Dec. 16-17, 1988 (Abst).
- Hastings, W.H. 1979 Nutritional requirements and feed technology. In: Advances in Aquaculture Fishing News Books Ltd., Farnham, Surrey, England (eds. Pillay, T.V.R. and Dill, Wm. A.), pp. 568-574.
- Hastings, W.H. and Dupree, H.K. 1969 Formula feeds for channel catfish. Prog. Fish Cult. 31: 187-196.
- Hilton, J.W. 1983 Potential of freeze dried worm meal as a replacement for fish meal in trout diet formulations. Aquaculture 32: 277-283.
- Jafri, A.K., Khawaja, D.K. and Qasim, S.Z. 1964 Studies on the biochemical composition of some freshwater fish. I. Muscles. Fish. Technol. 1: 148-157.
- Jauncey, K. 1982 Carp (Cyprinus carpio) Nutrition- A Review. In: Advances in Aquaculture Fishing News Books Ltd., Farnham, Surrey, England (eds. Pillay, T.V.R. and Dill, Wm. A.), pp. 215-263.

- Jhingran, V.G. 1983 Fish and Fisheries of India, Hindustan Publ. Co., New Delhi, India. 666 p.
- Jobling, M. and Wandsvik 1983 Quantitative protein requirements of Arctic charr, Salvelinus alpinus (L). J. Fish. Biol. 22: 705-712.
- Kalyani, R. and Shetty, H.P.C. 1987 Use of bio-gas slurry as a fertilizer for production of carps. In: The First Indian Fisheries Forum, Mangalore, Karnataka, Dec. 4-8, 1987 (Abst).
- Kanazawa, A. 1985 Essential fatty acid and lipid requirement of fish. In: Nutrition and Feeding in Fish. Academic Press, London/New York. (eds. Cowey, C.B., Mackie, A.M. and Bell, J.G.). pp. 281-298.
- Ketola, H.G. 1982 Amino acid nutrition of fishes. Requirements and supplementation of diets. Comp. Biochem. Physiol. 73B: 17-24.
- Kirchgessner, M., Kurzinger, H. and Schwarz, F.G. 1986 Digestibility of crude nutrients in different feeds and estimation of their energy content for carp (Cyprinus carpio L.). Aquaculture 58: 185-194.
- Klinnavee, S., Tansakul, R. and Promkuntong, V. 1989 Growth of nile tilapia (Oreochromis niloticus) fed with aquatic plant mixtures. In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst).
- Lakshmanan, M.A.V., 1967
Murty, D.S.,
Pillai, K.K. and
Banerjee, S.C. On a new artificial feed for carp fry. FAO. Fish. Rep. (44) 3: 373-87.
- Lall, S.P. 1979 Minerals in finfish nutrition. In: Finfish Nutrition and Fish-feed Technology. Vol. I Heenemann, Berlin, Federal Republic of Germany (eds. Halver, J.E. and Tiews, K.) pp. 85-97.

- | | | |
|---|-------|--|
| Law, A.T. | 1986 | Digestibility of low-cost ingredients in pelleted feed by grass carp (<u>Ctenopharyngodon idella</u> C. et V.) <u>Aquaculture</u> 51: 97-103. |
| Leary, D.F. and Lovell, R.T. | 1975 | Value of fibre in production type diet for channel catfish. <u>Trans. Am. Fish. Soc.</u> 104: 328-332. |
| Lee, D.J. and Putnam, G.B. | 1973 | The response of rainbow trout to varying protein/energy ratios in a test diet. <u>J. Nutr.</u> 103: 916-922. |
| Lovell, R.T. | 1976 | Voluntary food consumption by channel catfish offered expanded feeds of two nutrient densities. <u>Fisheries Ann. Rep., Auburn Agric. Expt. Sta., Auburn, AL.</u> pp. 1-31. |
| Lovell, R.T. | 1984a | Mineral requirements. In: <u>Nutrition and Feeding of Channel Catfish (Revised), South. Coop. Ser. Bull.</u> (eds. Robinson, E.H. and Lovell, R.T.). No. 296, pp. 26-28. |
| Lovell, R.T. | 1984b | Energy requirements. In: <u>Nutrition and Feeding of Channel Catfish (Revised), South. Coop. Ser. Bull.</u> (eds. Robinson, E.H. and Lovell, R.T.). No. 296, pp. 12-14. |
| Machiels, M.A.M. and Henken, A.M. | 1985 | Growth rate, feed utilization and energy metabolism of the African catfish, <u>Clarias gariepinus</u> (Burchell, 1822), as affected by dietary protein and energy content. <u>Aquaculture</u> 44: 271-284. |
| Maria, A.S., Hanifa, M.A. and Venkataramani, V.R. | 1987 | Studies on food utilisation in the common carp, <u>Cyprinus carpio</u> L. in relation to artificial feeds. In: <u>The First Indian Fisheries Forum. Mangalore, Karnataka, Dec. 4-8, 1987 (Abst).</u> |

- Marries, J.F.K. and Kissil, G.Wm. 1979 The influence of energy level on the feed intake, growth, food conversion and body composition of Sparus aurata. Aquaculture 17: 203-219.
- Mertz, E.T. 1972 The protein and amino acid needs. In: Fish Nutrition. Academic Press, London/New York. (ed. Halver, J.E.). pp. 106-143.
- Millikin, M.R. 1982 Qualitative and quantitative nutrient requirements of fishes; A Review. Fisheries Bulletin 80(4): 655-686.
- Millikin, M.R. 1983 Interactive effects of dietary protein and lipid on growth and protein utilization of age-0 striped bass. Trans. Am. Fish. Soc. 112: 185-193.
- Mohanty, S.N., Swamy, D.N. and Kumar, K. 1987 A note on the efficiency of a commercial diet in rohu fry. In: The First Asian Fisheries Forum, Mangalore, Karnataka, Dec. 4-8, 1987 (Abst).
- Morrison, F.B. 1961 Feeds and Feeding. The Morrison Publishing Company, Cluiton, Iowa. 1165 p.
- Naegel, L.C.A. 1989 Dried azolla as a feed ingredient for tilapias. In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst).
- Nandeesha, M.C., Keshavanath, P., Varghese, T.J., Shetty, H.P.C., Srikanth, G.K. and Basavaraja, N. 1988 Formulation and evaluation of agricultural waste-based carp diets. In: National Symposium on Aquaculture Productivity, New Delhi, Dec. 16-17, 1988 (Abst).



- Nandeesh, M.C., Srikant, G.K., Keshavanath, P., Varghese, T.J. and Shetty, H.P.C. 1989 Growth performance of Cyprinus carpio Var. communis fed on diets containing different levels of deoiled silkworm pupa. In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst).
- NAS-NRC 1981 Nutrient Requirements of Coldwater Fishes. No. 16, National Academy of Sciences Press, Washington D.C. 63 p.
- NAS-NRC 1983 Nutrient Requirements of Warmwater Fishes and Shellfishes, National Academy of Sciences Press, Washington D.C. 102 p.
- Naskar, R. and Sen, P.R. 1987 Utilization of duckweed for carp seed rearing. In: The First Indian Fisheries Forum, Mangalore, Karnataka, Dec. 4-8, 1987 (Abst).
- Niamat, R. 1982 Formulation of artificial feed and its nutritional efficiency test on catfish, Heteropneustes fossilis (Bloch). M.Phil. Dissertation, Aligarh Muslim University, Aligarh, India.
- Niamat, R. and Jafri, A.K. 1984a Preliminary observations on the use of water hyacinth (Eichhornia crassipes) leaf meal as protein source in fish feed. Curr. Sci. 53(6): 338-340.
- Niamat, R. and Jafri, A.K. 1984b Growth response of the siluroid, Heteropneustes fossilis Bloch, fed pelleted feed. Aquaculture 37: 223-229.
- Nose, T. and Arai, S. 1979 Recent advances in studies on mineral nutrition of fish in Japan. In: Advances in Aquaculture. Fishing News Books Ltd., Farnham, Surrey, England (eds. Pillay, T.V.R. and Dill, Wm. A.) pp. 584-590.

- Ogino, C. 1979 The present situation of studies of fish nutrition. In: Proceedings of the 7th Japan-Soviet Symposium on Aquaculture, Tokyo, Sept. 1978 pp. 11-18.
- Oser, L.B. 1965 In: Hawk's Physiological Chemistry (ed. Oser, L.B.) 14th. edition. The Blakiston Division Mcgraw. Hill. Beak Company, London. 1472 p.
- Otubusin, S.O. 1987 Effect of different levels of blood meal in pelleted feeds on tilapia, Oreochromis niloticus, production in floating bamboo net-cages. Aquaculture 65(3/4): 263-266.
- Page, J.W. and Andrews, J.W. 1973 Interactions of dietary levels of protein and energy on channel catfish. J. Nutr. 103: 1339-1346.
- Pandian, T.J. 1967 Intake, digestion and conversion of food in the fishes Megalops cyprinoides and Ophiocephalus striatus, Mar. Biol. 1: 16-32.
- Pandian, T.J. 1970 Intake and conversion of food in the fish Limonda limonda exposed to different temperatures. Mar. Biol. 5: 1-17.
- Pappathi, R. and Paul Raj, S. 1987 Fish feed formulations using wild leaves from Western Ghats. In: The First Indian Fisheries Forum, Mangalore, Karnataka, Dec. 4-8, 1987 (Abst).
- Patra, B.C., Das, I. and Ray, A.K. 1989 Nutritive value of aquatic weed Nymphoides cristatum (Roxb.) Q. kuntze as feed for Indian major carp, Labeo rohita (Hamilton). In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst).
- Paul Raj, S. 1989 Evaluation of Clitoria ternatia leaf as a protein supplement in the feed of Cyprinus carpio Var. communis. In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst).

- o Pieper, A. and Pfeffer, E. 1979 Carbohydrates as possible sources of dietary energy for rainbow trout (Salmo gairdneri, Richardson). In: Finfish Nutrition and Fishfeed Technology. Vol. I, Heenemann, Berlin, Federal Republic of Germany, pp. 209-219.
- Pillay , T.V.R. 1976 The state of Aquaculture, 1975. FAO Technical Conference on Aquaculture, Kyoto, Japan, May 26-June, 2, 1976. FIR/AQ/Conf./76/R.36.
- Pillay, T.V.R. 1986. Aquaculture; An International Perspective. In: Fisheries Development : 2000 AD. Oxford and IBH publishing Co., New Delhi/Bombay/Calcutta (ed. Trivedi K.K.). pp. 153-160.
- Prejs, A. and Blaszezyk, M. 1977 Relationships between food and cellulase activity in freshwater fishes. J. Fish. Biol. 11: 447-452.
- Rao, S.C., 1987 On the utility of selected organic manure as supplemented feed for the fingerlings of Catla catla (Hamilton). In: The First Indian Fisheries Forum Mangalore, Karnataka, Dec. 4-8, 1987 (Abst).
- Reinitz, G. and Hitzel, F. 1980 Formulation of practical diets for rainbow trout based on desired performance and body composition. Aquaculture 19: 243-252.
- Renukaradhya, K.M. and Varghese, T.J. 1986 Protein requirements of the carps Catla catla (Hamilton) and Labeo rohita (Hamilton). Proc. Indian Acad. Sci. 95 (1): 103-107.
- Ringrose, R.C. 1971 Calorie-to-protein ratio for brook trout (Salvelinus fontinalis) J. Fish. Res. Bd. Can. 28: 1113-1117.

- Robinson, E.H. 1984 Vitamin requirements. In: Nutrition and Feeding of Channel Catfish (Revised). South. Coop. Ser. Bull. (eds. Robinson, E.H. and Lovell R.T.). No. 296, pp. 21-25.
- Rozin, P. and Mayer, J. 1961 Regulation of food intake in the goldfish. Am. J. Physiol. 201: 968-974.
- Sen, P.R., Rao, N.G.S., Ghosh, S.R. and Rout, M. 1978 Observations on the protein and carbohydrate requirements of carps. Aquaculture 13: 245-255.
- Shimeno, S., Hosokawa, H., Takeda, M. and Kojiyama, H. 1980 Effects of calorie to protein ratios in formulated diet on the growth, feed conversion and body composition of young yellow tail. Bull. Jap. Soc. scient. Fish. 46: 1083-1087.
- Sinha, V.R.P. 1988 Aquaculture Research and Development in India. In: International Symposium on Aquaculture Research Needs for the year 2000, New Delhi, India, Nov. 15-18, 1988.
- Spinelli, J., Mahnken, C. and Steinberg, M. 1979 Alternate sources of proteins for fish meal in salmonid diets. In: Finfish Nutrition and Fish-feed Technology Vol. II. Heene-mann, Berlin, Federal Republic of Germany (eds. Halver, J.E. and Tiews, K.). pp. 131-142.
- Stickney, R.R. 1979 Feeds, nutrition and growth. In: Principles of Warmwater Aquaculture. John Wiley, Toronto, Ont. pp. 198-202.
- Stickney, R.R. 1984 Lipids. In: Nutrition and Feeding of Channel Catfish, South. Coop. Ser. Bull. (eds. Stickney, R.R. and Lovell, R.T.). No. 296 pp. 17-20.

- | | | |
|---|-------|---|
| Stickney, R.R. and Lovell, R.T. | 1977 | Nutrition and Feeding of Channel Catfish. <u>South. Coop. Ser. Bull.</u> No. 218: 67 pp. |
| Stickney, R.R. and Shumway, S.E. | 1974 | Occurrence of cellulase activity in the stomachs of fishes <u>J. Fish. Biol.</u> 6: 779-790. |
| Strange, R.J. | 1984 | Carbohydrates. In: Nutrition and Feeding of Channel Catfish, <u>South. Coop. Ser. Bull.</u> (eds. Stickney, R.R. and Lovell, R.T.). No. 296: pp. 15-16. |
| Sumagaysay, N.S., and Chiu, Y.N. | 1989 | Effects of dietary levels of energy and fibre on protein utilization of supplementary feeds of milkfish (<u>Chanos chanos</u> Forsskal) reared in brackishwater ponds. In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst). |
| o TAC | 1973 | Report of the TAC Working Groups on Aquaculture Consultative Group on International Agricultural Research. Technical Advisory Committee. 7th Meeting, Rome, Feb. 4-8, 1973. FAO. DDDR: IAR 73 (75) Nov.1973. |
| Tacon, A.G.J. and Jackson, A.J. | 1985 | Utilisation of conventional and unconventional protein sources in practical fish feeds. In: Nutrition and Feeding in Fish. Academic Press. London/New York. (eds. Cowey, C.B., Mackie, A.M. and Bell J.G.). pp. 119-145. |
| Takeuchi, T., Watanabe, T. and Ogino, C. | 1978a | Supplementary effect of lipids in a high protein diet of rainbow trout. <u>Bull. Jap. Soc. scient. Fish.</u> 44: 677-681. |
| Takeuchi, T., Watanabe, T., and Ogino, C. | 1978b | Optimum ratio of protein to lipid in diets of rainbow trout. <u>Bull. Jap. Soc. scient. Fish.</u> 44: 683-688. |

- Takeuchi, T.,
Watanabe, T., and
Ogino, C. 1978c Studies on nutritive value of dietary lipids in fish. XV. Use of hydrogenated fish oil and beef tallow as a dietary energy source for carp and rainbow trout, Bull. Jap. Soc. scient. Fish. 44: 875-881.
- Takeuchi, T.,
Watanabe, T., and
Ogino, C. 1978d Optimum ratio of energy to protein for carp, Bull. Jap. Soc. scient. Fish. 45: 983-987.
- Takeuchi, T.,
Watanabe, T., and
Ogino, C. 1979 Studies on nutritive value of dietary lipids in fish XVI. Availability of carbohydrates and lipids as dietary energy source for carp. Bull. Jap. Soc. scient. Fish. 45: 977-982.
- Takeuchi, T.,
Sato, S.S. and
Watanabe, T. 1983 Requirement of Tilapia nilotica for essential fatty acids. Bull. Jap. Soc. scient. Fish. 49: 1127-1137.
- Toor, H.S.,
Kaur, K. and
Dhawan, A. 1988 Utilization of biological wastes in carp culture. In: National Symposium on Aquaculture Productivity, New Delhi, Dec. 16-17, 1988 (Abst).
- Venkatesh, B.,
Mukherji, A.P. and
Dehadrai, P.V. 1986 Growth and metabolism of the catfish Clarias batrachus (Linn.) fed with different experimental diets. Proc. Indian Acad. Sci. 95(4): 457-462.
- Viola, S.,
Mokady, S.,
Rappaport, U. and
Arieli, Y. 1982 Partial and complete replacement of fishmeal by soybean meal in feeds for intensive culture of carp. Aquaculture 26: 223-236.
- Watanabe, T. 1977 Sparing action of lipids on dietary protein in fish - low protein diet with high calorie content. Technocrat 10: 34-39.
- Watanabe, T. 1982 Lipid nutrition in fish. Comp. Biochem. Physiol. 73B (1): 3-15.

- Watanabe, T. 1985 Importance of the study of broodstock nutrition for further development of aquaculture. In: Nutrition and Feeding in Fish. Academic Press, London/New York (eds. Cowey, C.B., Mackie, A.M. and Bell J.G.). pp. 395-414.
- Wee, L.L. and Little, D. 1989 Supplementary feeds for nile tilapia culture. In: The Second Asian Fisheries Forum, Tokyo, Japan, April 17-22, 1989 (Abst).
- Wilson, R.P. and Halver, J.E. 1986 Protein and amino acid requirements of fishes. Ann. Rev. Nutr. 6: 225-244.
- Yu, T.C. and Sinnhuber, R.O. 1981 Use of beef tallow as an energy source in coho salmon (Oncorhynchus kisutch) rations. Can. J. Fish. Aquat. Sci. 38: 367-370.